



ISPRA
Institute for Environmental
Protection and Research

Italian Emission Inventory 1990-2007

Informative Inventory Report 2009

Daniela Romano, Antonella Bernetti, Rocío D. Córdor,
Riccardo De Lauretis, Eleonora Di Cristofaro, Andrea Gagna,
Barbara Gonella, Ernesto Taurino, Marina Vitullo

ISPRA - Institute for Environmental Protection and Research

*Annual Report for submission under the UNECE Convention
on Long-range Transboundary Air Pollution*

May 2009

Legal Disclaimer

The Institute for Environmental Protection and Research, or persons acting on its behalf, are not responsible for the use that may be made of the information contained in this report.

Authors

Daniela Romano, Antonella Bernetti, Rocío D. Córdor,
Riccardo De Lauretis, Eleonora Di Cristofaro, Andrea Gagna,
Barbara Gonella, Ernesto Taurino, Marina Vitullo

Contact: Riccardo De Lauretis
telephone +39 0650072543
fax +39 0650072657
e-mail riccardo.delauritis@isprambiente.it

ISPRA- Institute for Environmental Protection and Research
Environment Department
Monitoring and Prevention of Atmospheric Impacts
Air Emission Inventory Unit
Via V. Brancati, 48 00144 Rome ITALY

ISPRA is the Institute for Environmental Protection and Research established by the Italian Law 133/2008, as published in the Official Journal n. 195, August 21 2008. The Institute performs the functions of three former institutions: APAT (Agency for Environmental Protection and Technical Services), ICRAM (Central Institute for Applied Marine Research), INFS (National Institute for Wildlife).

This publication refers to activities carried out prior to the unification of the three institutions and, therefore, individual reference is still made to them.

Extracts from this document may be reproduced on the condition that the source is acknowledged.

© ISPRA 2009

Contents

EXECUTIVE SUMMARY	5
1 INTRODUCTION	6
1.1 BACKGROUND INFORMATION ON THE CONVENTION ON LONG-RANGE TRANSBOUNDARY AIR POLLUTION	6
1.2 NATIONAL INVENTORY	7
1.3 INSTITUTIONAL ARRANGEMENTS	8
1.4 INVENTORY PREPARATION PROCESS	10
1.5 METHODS AND DATA SOURCES	11
1.6 KEY CATEGORIES	13
1.7 QA/QC AND VERIFICATION METHODS	16
1.8 GENERAL UNCERTAINTY EVALUATION	18
1.9 GENERAL ASSESSMENT OF COMPLETENESS	18
2 ANALYSIS OF KEY TRENDS BY POLLUTANT.....	19
2.1 MAIN POLLUTANTS	19
2.1.1 SULPHUR DIOXIDE (SO _x)	19
2.1.2 NITROGEN OXIDES (NO _x)	21
2.1.3 AMMONIA (NH ₃)	25
2.1.4 NON METHANE VOLATILE ORGANIC COMPOUNDS (NMVOC)	27
2.1.5 CARBON MONOXIDE (CO)	30
2.2 PARTICULATE MATTER	32
2.2.1 PM ₁₀	32
2.2.2 PM _{2.5}	34
2.3 HEAVY METALS (Pb, Cd, Hg)	36
2.3.1 LEAD (Pb)	36
2.3.2 CADMIUM (Cd).....	38
2.3.3 MERCURY (Hg)	40
2.4 PERSISTENT ORGANIC POLLUTANTS (POPs).....	42
2.4.1 POLYCYCLIC AROMATIC HYDROCARBONS (PAH).....	42
2.4.2 DIOXINS.....	44
2.4.3 HEXACHLOROBENZENE (HCB).....	45
2.4.4 POLYCHLORINATED BIPHENYL (PCB).....	47
3 ANALYSIS OF KEY TRENDS BY SECTOR.....	49
3.1 ENERGY (NFR SECTOR 1).....	49
3.1.1 METHODOLOGICAL ISSUES.....	49
3.2 INDUSTRIAL PROCESSES (NFR SECTOR 2).....	52
3.2.1 METHODOLOGICAL ISSUES.....	52
3.3 SOLVENT AND OTHER PRODUCT USE (NFR SECTOR 3).....	53
3.3.1 METHODOLOGICAL ISSUES.....	53
3.4 AGRICULTURE (NFR SECTOR 4)	55

3.4.1	METHODOLOGICAL ISSUES	55
3.5	WASTE (NFR SECTOR 6).....	57
3.5.1	METHODOLOGICAL ISSUES	57
4	RECALCULATIONS AND IMPROVEMENTS.....	59
4.1	RECALCULATIONS	59
4.2	PLANNED IMPROVEMENTS	61
5	PROJECTIONS.....	62
	REFERENCES	65
	ANNEX: NFR CODES.....	68

EXECUTIVE SUMMARY

This report is the second Italian Informative Inventory Report (IIR) in the framework of the United Nations Economic Commission for Europe (UNECE) Convention on Long Range Transboundary Air Pollution (CLRTAP). It contains information on the Italian inventory up to the year 2007, including an explanation of methodologies, data sources, QA/QC activities and verification processes carried out during the inventory compilation, with an analysis of emission trends and a description of key categories.

The aim of the document is to facilitate understanding of the calculation of the Italian air pollutant emission data, hence providing a common means for comparing the relative contribution of different emission sources and helping in the identification of reduction policies.

The Institute for Environmental Protection and Research (ISPRA) has the overall responsibility for the emission inventory submission to CLRTAP, as well as to the United Nations Framework Convention on Climate Change (UNFCCC), and is in charge of all work related to inventory compilation.

In particular, in compliance with the LRTAP Convention, Italy has to submit annually data on national emissions of SO_x, NO_x, NMVOC, CO and NH₃, and various heavy metals and POPs. The submission consists of the national emission inventory, communicated through compilation of the Nomenclature Reporting Format (NRF), and the informative inventory report (IIR) to ensure the properties of transparency, consistency, comparability, completeness and accuracy.

In the period 1990-2007, emissions from most pollutants described in this report show a downward trend. Reductions are especially relevant for the main pollutants (SO_x -81%; NO_x -43%; CO -52%; NMVOC -38%) and lead (-94%) whereas a significant raise is observed for polycyclic aromatic hydrocarbons (+51%).

The major drivers for the trend are reductions in the industrial and road transport sectors, due to the implementation of various European Directives which introduced new technologies, plant emission limits, the limitation of sulphur content in liquid fuels and the shift to cleaner fuels. Emissions have also decreased for the improvement of energy efficiency as well as the promotion of renewable energy.

The energy sector is the main source of emissions in Italy with a share of more than 80% in many pollutants (SO_x 80%; NO_x 98%; CO 88%; PM_{2.5} 81%). The industrial processes sector is an important source of emissions specifically related to the iron and steel production, at least for particulate matter, heavy metals and POPs, whereas significant emissions of SO_x and particulate matter derive from cement production; on the other hand, the solvent and other product use sector is characterized by NMVOC emissions. The agriculture sector is the main source of NH₃ emissions in Italy with a share of 95% in national total. Finally, the waste sector, specifically waste incineration, is a relevant source for HCB, PAH and dioxin emissions (50%, 20% and 12%, respectively).

Emission figures of the Italian emission inventory and other related documents are publicly available at http://www.sinanet.apat.it/it/sinanet/serie_storiche_emissioni.

1 INTRODUCTION

1.1 Background information on the Convention on Long-range Transboundary Air Pollution

The 1979 Geneva Convention on Long-range Transboundary Air Pollution, contributing to the development of international environmental law, is one of the fundamental international means for the protection of the human health and the environment through the intergovernmental cooperation. The fact that air pollutants could travel several thousands of kilometres before deposition and damage occurred outlined the need for international cooperation.

In November 1979, in Geneva, 34 Governments and the European Community (EC) signed the Convention. The Convention on Long-range Transboundary Air Pollution was ratified by Italy in the year 1982 and entered into force in 1983. It has been extended by the following eight specific protocols:

- The 1984 Protocol on Long-term Financing of the Cooperative Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe (EMEP); 42 Parties. Entered into force on 28th January 1988.
- The 1985 Protocol on the Reduction of Sulphur Emissions or their Transboundary Fluxes by at least 30 per cent; 23 Parties. Entered into force on 2nd September 1987.
- The 1988 Protocol concerning the Control of Nitrogen Oxides or their Transboundary Fluxes; 31 Parties. Entered into force on 14th February 1991.
- The 1991 Protocol concerning the Control of Emissions of Volatile Organic Compounds or their Transboundary Fluxes; 22 Parties. Entered into force on 29th September 1997.
- The 1994 Protocol on Further Reduction of Sulphur Emissions; 27 Parties. Entered into force on 5th August 1998.
- The 1998 Protocol on Heavy Metals; 28 Parties. Entered into force on 29 December 2003.
- The 1998 Protocol on Persistent Organic Pollutants (POPs); 28 Parties. Entered into force on 23rd October 2003.
- The 1999 Protocol to Abate Acidification, Eutrophication and Ground-level Ozone; 23 Parties. Entered into force on 17th May 2005. (Guidance documents to Protocol adopted by decision 1999/1).

As regards Italy, the following table shows the dates of signature and ratification of both Convention and Protocols.

	Signature	Ratification
1979 Convention	14/11/1979	15/07/1982
1984 EMEP Protocol	28/09/1984	12/01/1989
1985 Sulphur Protocol	09/07/1985	05/02/1990
1988 NO _x Protocol	01/11/1988	19/05/1992
1991 VOC Protocol	19/11/1991	30/06/1995
1994 Sulphur Protocol	14/06/1994	14/09/1998
1998 Heavy Metals Protocol	24/06/1998	
1998 POPs Protocol	24/06/1998	20/06/2006
1999 Multi-effect Protocol	01/12/1999	

Table 1.1 Dates of signature and ratification of the UNECE Convention and Protocols

The following classes of pollutants should be included in the emission inventory:

Main Pollutants

- Sulphur oxides (SO_x), in mass of SO₂;
- Nitrous oxides (NO_x), in mass of NO₂;
- Carbon monoxide (CO);
- Non-methane volatile organic compounds (NMVOC);
- Ammonia (NH₃).

Particulate matter

- PM10, particulate matter less than 10 microns in diameter;
- PM2.5, particulate matter less than 2.5 microns in diameter.

Heavy Metals

- Priority Metals: Lead (Pb), Cadmium (Cd) and Mercury (Hg);
- Other metals: Arsenic (As), Chrome (Cr), Copper (Cu), Nickel (Ni), Selenium (Se) and Zinc (Zn).

Persistent organic pollutants (POPs)

- As specified in Annex I of the POPs Protocol;
- As specified in Annex II of the POPs Protocol, including Polychlorinated Biphenyls (PCBs);
- As specified in Annex III of the POPs Protocol: Dioxins (Diox), Polycyclic Aromatic Hydrocarbons (PAHs), Hexachlorobenzene (HCB);
- Other POPs.

1.2 National Inventory

As a Party to the United Nations Economic Commission for Europe (UNECE) Convention on Long Range Transboundary Air Pollution (CLRTAP), Italy has to submit annually data on emissions of air pollutants in order to fulfil obligations, in compliance with the implementation of Protocols under the Convention. Parties are required to report on annual national emissions of SO_x, NO_x, NMVOC, CO and NH₃, and various heavy metals and POPs according to the Guidelines for Estimating and Reporting Emission Data under the Convention on Long-range Transboundary Air Pollution (UNECE, 2003).

Specifically, the submission consists of the national LRTAP emission inventory, communicated through compilation of the Nomenclature Reporting Format (NRF) and the informative inventory report (IIR).

The Italian informative inventory report contains information on the national inventory for the year 2006, including descriptions of methods, data sources, QA/QC activities carried out and a trend analysis. The inventory accounts for anthropogenic emissions of the following substances: sulphur oxides (SO_x), nitrogen oxides (NO_x), ammonia (NH₃), non-methane volatile organic compounds (NMVOC), carbon monoxide (CO), particulate matter, particles of size <10 µm, (PM10), particulate matter, particles of size < 2.5µm, (PM2.5), lead (Pb), cadmium (Cd), mercury (Hg), arsenic (As), chromium (Cr), copper (Cu), nickel (Ni), selenium (Se), zinc (Zn), polychlorinated biphenyls (PCBs), polycyclic aromatic hydrocarbons (PAH), dioxins (Diox), hexachlorobenzene (HCB). Other pollutants are reported either as not estimated or not occurring, further investigation is planned to verify these emissions.

Detailed information on emission figures of primary pollutants, particulate matter, heavy metals and persistent organic pollutants as well as estimation procedures are provided in order to improve the transparency, consistency, comparability, accuracy and completeness of the inventory provided.

The national inventory is updated annually in order to reflect revisions and improvements in the methodology and the availability of new information. Adjustments are applied retrospectively to

earlier years, which accounts for any difference in previously published data.
Total emissions from 1990 to 2007 are reported in Table 1.2 by pollutant.

		1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
SO_x	<i>Gg</i>	1,795	1,319	749	697	616	518	480	401	379	338
NO_x	<i>Gg</i>	2,003	1,864	1,428	1,416	1,361	1,354	1,314	1,224	1,182	1,141
NMVOC	<i>Gg</i>	1,939	2,002	1,566	1,501	1,432	1,374	1,320	1,249	1,222	1,195
NH₃	<i>Gg</i>	466	448	451	452	439	434	426	414	410	418
CO	<i>Gg</i>	6,928	6,877	4,860	4,648	4,221	4,067	3,884	3,509	3,345	3,336
As	<i>Mg</i>	37	27	45	45	41	42	41	40	41	41
Cd	<i>Mg</i>	10	10	9	9	7	8	8	8	9	9
Cr	<i>Mg</i>	89	70	48	48	48	51	53	55	56	56
Cu	<i>Mg</i>	98	104	102	102	102	104	107	108	110	110
Hg	<i>Mg</i>	12	11	10	10	10	10	10	10	11	11
Ni	<i>Mg</i>	122	113	104	109	112	113	111	110	108	104
Pb	<i>Mg</i>	4,371	1,988	935	702	236	241	256	265	273	273
Se	<i>Mg</i>	10	10	11	11	11	12	12	12	12	12
Zn	<i>Mg</i>	897	874	835	837	836	853	900	906	977	970
PM10	<i>Gg</i>	232	229	192	191	179	175	177	165	162	162
PM2.5	<i>Gg</i>	199	195	161	158	147	143	144	133	130	130
PAH	<i>Mg</i>	103	121	128	129	118	123	141	138	143	155
Dioxin	<i>g IT_q</i>	472	460	369	293	283	282	289	294	302	318
HCB	<i>Mg</i>	0.019	0.020	0.021	0.030	0.032	0.033	0.024	0.021	0.028	0.028
PCB	<i>Mg</i>	0.242	0.252	0.226	0.231	0.237	0.240	0.245	0.237	0.246	0.237

Table 1.2 Emission time series by pollutant

The NRF files and other related documents can be found on website at the following address:
http://www.sinanet.apat.it/it/sinanet/serie_storiche_emissioni.

1.3 Institutional arrangements

The Institute for Environmental Protection and Research (ISPRA) has the overall responsibility for the emission inventory and submissions to CLRTAP; the institute is also responsible for the communication of the pollutants under the NEC directive as well as to carry out scenarios, jointly with the Agency for Energy, New technologies and Environment (ENEA), as established by the

Legislative Decree n. 171 of 21st May 2004. Moreover, ISPRA (former APAT) is the single entity in charge of the development and compilation of the national greenhouse gas emission inventory as indicated by the Legislative Decree n. 51 of 7th March 2008. The Ministry for the Environment, Land and Sea is responsible for the endorsement of the inventory and for the communication to the Secretariat of the different conventions.

The Italian National System, currently in place, is fully described in the document 'National Greenhouse Gas Inventory System in Italy' (ISPRA, 2009 [a]).

A specific unit of the Institute is responsible for the compilation of the Italian Atmospheric Emission Inventory and the Italian Greenhouse Gas Inventory in the framework of both the Convention on Climate Change and the Convention on Long Range Transboundary Air Pollution. The whole inventory is compiled by the institute; scientific and technical institutions and consultants may help in improving information both on activity data and emission factors of specific activities. All the measures to guarantee and improve the transparency, consistency, comparability, accuracy and completeness of the inventory are undertaken.

ISPRA bears the responsibility for the general administration of the inventory, co-ordinates participation in review processes, publishes and archives the inventory results.

Specifically, ISPRA is responsible for all aspects of national inventory preparation, reporting and quality management. Activities include the collection and processing of data from different data sources, the selection of appropriate emissions factors and estimation methods consistent with the EMEP/CORINAIR guidelines, the IPCC 1996 Revised Guidelines, the IPCC Good Practice Guidance and Uncertainty management and the IPCC Good Practice Guidance for land use, land-use change and forestry, and the IPCC 2006 Guidelines, the compilation of the inventory following the QA/QC procedures, the preparation of the Informative Inventory Report and the reporting through the Nomenclature Reporting Format, the response to review checks, the updating and data storage.

Different institutions are responsible for statistical basic data and data publication, which are primary to ISPRA for carrying out emission estimates. These institutions are part of the National Statistical System (Sistan), which provides national official statistics, and therefore are asked periodically to update statistics; moreover, the National Statistical System ensures the homogeneity of the methods used for official statistics data through a coordination plan, involving the entire public administration at central, regional and local levels.

The main Sistan products, which are primarily necessary for the inventory compilation, are:

- National Statistical Yearbooks, Monthly Statistical Bulletins, by ISTAT (National Institute of Statistics);
- Annual Report on the Energy and Environment, by ENEA (Agency for New Technologies, Energy and the Environment);
- National Energy Balance (annual), Petrochemical Bulletin (quarterly publication), by MSE (Ministry of Economic Development);
- Transport Statistics Yearbooks, by MINT (Ministry of Transportation);
- Annual Statistics on Electrical Energy in Italy, by TERNA (National Independent System Operator);
- Annual Report on Waste, by ISPRA (former APAT);
- National Forestry Inventory, by MIPAAF (Ministry of Agriculture, Food and Forest Policies).

The national emission inventory itself is a Sistan product.

Other information and data sources are used to carry out emission estimates, which are generally referred to in Table 1.3 in the following section 1.5.

1.4 Inventory preparation process

ISPRA has established fruitful cooperation with a number of governmental and research institutions as well as industrial associations, which helps improving some leading categories of the inventory. Specifically, these activities aim at the improvement of provision and collection of basic data and emission factors, through plant-specific data, and exchange of information on scientific researches and new sources. Moreover, when in depth investigation is needed and a high uncertainty in the estimates is present, specific sector analyses are committed to ad hoc research teams or consultants. ISPRA also coordinates with different national and regional authorities and private institutions for the cross-checking of parameters and estimates, as well as with ad hoc expert panels, in order to improve the completeness and transparency of the inventory.

The main basic data needed for the preparation of the national emission inventory are energy statistics, published by the Ministry of Economic Development (MSE) in the National Energy Balance (BEN), statistics on industrial and agricultural production, published by the National Institute of Statistics (ISTAT), statistics on transportation, provided by the Ministry of Transportation (MINT), and data supplied directly by the relevant professional associations.

Emission factors and methodologies used in the estimation process are consistent with the EMEP/CORINAIR Guidebook, the IPCC Guidelines and Good Practice Guidance as well as supported by national experiences and circumstances. Final decisions are up to inventory experts, taking into account all the information available.

For the industrial sector, emission data collected through the National Pollutant Emission Register (EPER, now E-PRTR), the Large Combustion Plant (LCP) Directive and in the framework of the European Emissions Trading Scheme have yielded considerable developments in the inventory of the relative sectors. In fact, these data, even if not always directly used, are taken into account as a verification of emission estimates and improve national emissions factors as well as activity data figures.

In addition, final estimates are checked and verified also in view of annual environmental reports by industries.

For large industrial point sources, emissions are registered individually, when communicated, based upon detailed information such as fuel consumption.

Other small plants communicate their emissions which are also considered individually.

Emission estimates are drawn up for each sector. Final data are communicated to the UNECE Secretariat filling in the NRF files.

The process of the inventory preparation is carried out annually. In addition to a new year, the entire time series is checked and revised during the annual compilation of the inventory. In particular, recalculations are elaborated on account of changes in the methodologies used to carry out emission estimates, changes due to different allocation of emissions as compared to previous submissions and changes due to error corrections. The inventory may also be expanded by including categories not previously estimated if sufficient information on activity data and suitable emission factors have been identified and collected. Information on the major recalculations is provided in the sectoral chapter of the report.

All the reference material, estimates and calculation sheets, as well as the documentation on scientific papers and the basic data needed for the inventory compilation, are stored and archived at the Institute. After each reporting cycle, all database files, spreadsheets and electronic documents are archived as 'read-only-files' so that the documentation and estimates could be traced back during the review process or the new year inventory compilation.

Technical reports and emission figures are publicly accessible on the web at the address http://www.sinanet.apat.it/it/sinanet/serie_storiche_emissioni.

1.5 Methods and data sources

An outline of methodologies and data sources used in the preparation of the emission inventory for each sector is provided in the following. In Table 1.3 a summary of the activity data and sources used in the inventory compilation is reported.

Methodologies are consistent with the EMEP-CORINAIR Emission Inventory Guidebook, Revised 1996 and 2006 IPCC Guidelines, and IPCC Good Practice Guidance (EMEP/CORINAIR, 2005; IPCC, 1997; IPCC, 2006; IPCC, 2000); national emission factors are used as well as default emission factors from international guidebooks, when national data are not available. The development of national methodologies is supported by background documents.

The most complete document describing national methodologies used in the emission inventory compilation is the National Inventory Report, submitted in the framework of the UN Convention on Climate Change and the Kyoto Protocol (ISPRA, 2009 [b]).

SECTOR	ACTIVITY DATA	SOURCE
1 Energy 1A1 Energy Industries	Fuel use	Energy Balance - Ministry of Economic Development Major national electricity producers European Emissions Trading Scheme
1A2 Manufacturing Industries and Construction	Fuel use	Energy Balance - Ministry of Economic Development Major National Industry Corporation European Emissions Trading Scheme
1A3 Transport	Fuel use Number of vehicles Aircraft landing and take-off cycles and maritime activities	Energy Balance - Ministry of Economic Development Statistical Yearbooks - National Statistical System Statistical Yearbooks - Ministry of Transportation Statistical Yearbooks - Italian Civil Aviation Authority (ENAC) Maritime and Airport local authorities
1A4 Residential-public-commercial sector	Fuel use	Energy Balance - Ministry of Economic Development
1B Fugitive Emissions from Fuel	Amount of fuel treated, stored, distributed	Energy Balance - Ministry of Economic Development Statistical Yearbooks - Ministry of Transportation Major National Industry Corporation
2 Industrial Processes	Production data	National Statistical Yearbooks- National Institute of Statistics International Statistical Yearbooks-UN European Emissions Trading Scheme European Pollutant Emission Registry Sectoral Industrial Associations
3 Solvent and Other Product Use	Amount of solvent use	National Environmental Publications - Sectoral Industrial Associations International Statistical Yearbooks - UN
4 Agriculture	Agricultural surfaces Production data Number of animals Fertiliser consumption	Agriculture Statistical Yearbooks - National Institute of Statistics Sectoral Agriculture Associations
5 Land Use, Land Use Change and Forestry	Forest and soil surfaces Amount of biomass Biomass burnt Biomass growth	Statistical Yearbooks - National Institute of Statistics State Forestry Corps National and Regional Forestry Inventory Universities and Research Institutes
6 Waste	Amount of waste	National Waste Cadastre - Institute for Environmental Protection and Research , National Waste Observatory

Table 1.3 Main activity data and sources for the Italian Emission Inventory

Activity data used in emission calculations and their sources are briefly described herebelow.

In general, for the energy sector, basic statistics for estimating emissions are fuel consumption published in the national Energy Balance by the Ministry of Economic Development. Additional information for electricity production is provided by the major national electricity producers and by the major national industry corporation. On the other hand, basic information for road transport, maritime and aviation, such as the number of vehicles, harbour statistics and aircraft landing and take-off cycles are provided in statistical yearbooks published both by the National Institute of Statistics and the Ministry of Transportation. Other data are communicated by different category associations.

The analysis of data from the Italian Emissions Trading Scheme database is used to develop country-specific emission factors and check activity data levels. In fact, ISPRA (former APAT) is also responsible for developing, operating and maintaining the national registry under Directive 2003/87/CE as instituted by the Legislative Decree 51 of March 7th 2008; the Institute performs this

tasks under the supervision of the national Competent Authority for the implementation of directive 2003/87/CE, jointly established by the Ministry for Environment, Land and Sea and the Ministry for Economic Development.

For the industrial sector, the annual production data are provided by national and international statistical yearbooks. Emission data collected through the National Pollutant Emission Register (EPER, now E-PRTR) are also used in the development of emission estimates or taken into account as a verification of emission estimates for some specific categories. According to the Italian Decree of 23 November 2001, data from the Italian E-PRTR are validated and communicated by ISPRA to the Ministry for the Environment, Land and Sea and to the European Commission within October of the current year for data referring to the previous year. These data are used for the compilation of the inventory whenever they are complete in terms of sectoral information; in fact, industries communicate figures only if they exceed specific thresholds; furthermore, basic data such as fuel consumption are not supplied and production data are not split by product but reported as an overall value. Anyway, E-PRTR is a good basis for data checks and a way to facilitate contacts with industries which, in many cases, supply, under request, additional information as necessary for carrying out sectoral emission estimates.

In addition, final emissions are checked and verified also taking into account figures reported by industries in their annual environmental reports.

Both for energy and industrial processes, emissions of large industrial point sources are registered individually; communication also takes place in the framework of the European Directive on Large Combustion Plants, based upon detailed information such as fuel consumption. Other small plants communicate their emissions which are also considered individually.

For the other sectors, i.e. for solvents, the amount of solvent use is provided by environmental publications of sector industries and specific associations as well as international statistics.

For agriculture, annual production data and number of animals are provided by the National Institute of Statistics and other sectoral associations.

For land use, land use change and forestry, forest and soil surfaces are provided by the National Institute of Statistics while statistics on forest fires are supplied by the State Forestry Corps.

For waste, the main activity data are provided by the Institute for Environmental Protection and Research and the Waste Observatory.

In case basic data are not available proxy variables are considered; unpublished data are used only if supported by personal communication and confidentiality of data is respected.

All the material and documents used for the inventory emission estimates are stored at the Institute for Environmental Protection and Research. The inventory is composed by spreadsheets to calculate emission estimates; activity data and emission factors as well as methodologies are referenced to their data sources.

A 'reference' database has also been developed to increase the transparency of the inventory; at the moment, it is complete as far as references to greenhouse gas emissions are concerned.

1.6 Key categories

A key category analysis of the Italian inventory is carried out according to the Tier 1 method described in the IPCC Good Practice (IPCC, 2000). According to these guidelines, a key category is defined as an emission category that has a significant influence on a country's inventory in terms of the absolute level in emissions. Key categories are those which, when summed together in descending order of magnitude, add up to over 95% of the total emissions.

National emissions have been disaggregated into the categories reported in the National Format Report; details vary according to different pollutants in order to reflect specific national circumstances. The level analysis has been applied to the last submitted inventory, as for 2007. Results are reported in the following table by pollutant.

	Key categories											Total (%)	
SO_x	1A1a (24.1%)	1A2 (19.3%)	1A1b (14.3%)	1A3d ii (13.4%)	1B2 (9.4%)	2A1 (4.2%)	1A1c (4.0%)	1B2c (2.7%)	1A4b (2.5%)	2B (2.4%)			96.2
NO_x	1A3b iii (21.9%)	1A3b i (21.0%)	1A2 (14.8%)	1A3 d ii (7.5%)	1A3b ii (6.9%)	1A4c (6.5%)	1A1a (6.0%)	1A4b (4.2%)	1A4a (3.5%)	1A1b (2.2%)	6C (1.2%)		95.6
NH₃	4D1 (40.1%)	4B1b (17.0%)	4B1a (15.8%)	4B8 (8.9%)	4B9 (7.4%)	1A3b i (3.3%)	4B13 (2.7%)						95.1
NMVOC	3A (18.8%)	1A3b iv (16.4%)	3D (15.9%)	1A3d ii (8.1%)	1A3b i (6.7%)	3C (6.6%)	1A4b (4.1%)	2D2 (2.3%)	1B2a iv (2.2%)	1B2b (2.1%)	1A3b v (2.0%)	3B (1.8%)	95.5
	1A4c (1.8%)	1B2a v (1.7%)	1A3b iii (1.4%)	1A4a (1.3%)	6C (1.2%)	1A2 (1.0%)							
CO	1A3b i (23.3%)	1A3b iv (19.2%)	1A4b (17.2%)	1A2 (11.1%)	6C (8.1%)	1A3d ii (5.8%)	1A4c (3.71%)	2C (3.2%)	1A3b ii (2.3%)	1A3b iii (1.7%)			95.6
PM10	1A2 (14.8%)	1A4b (14.5%)	6C (7.4%)	1A4c (7.1%)	1A3b i (6.9%)	1A3b vi (6.0%)	4B9 (5.7%)	1A3b ii (4.7%)	2C (4.6%)	1A3b iii (4.6%)	1A3d ii (3.9%)	2A1 (3.8%)	95.4
	4B8 (2.2%)	1A3b iv (1.9%)	2A6 (1.6%)	1A1a (1.4%)	4F (1.3%)	2A2 (1.2%)	4B1b (0.8%)	1A5b (0.8%)					
PM2.5	1A2 (17.6%)	1A4b (17.1%)	1A4c (8.9%)	1A3b i (8.6%)	6C (7.9%)	1A3b ii (5.8%)	1A3b iii (5.7%)	1A3d ii (4.9%)	2C (4.5%)	1A3b vi (4.1%)	1A3b iv (2.4%)	1A1a (1.7%)	95.1
	4F (1.7%)	1A5b (0.9%)	4B9 (0.9%)	1A4a (0.9%)	1A1b (0.8%)	2A1 (0.7%)							
Pb	1A2 (52.2%)	2C (28.8%)	1A4a (14.2%)										95.2
Cd	1A2 (38.0%)	1A4a (24.2%)	2C (15.1%)	1A4b (9.6%)	1B1b (2.7%)	1A3b i (2.6%)	6C (2.0%)	1A1a (1.5%)					95.6
Hg	1A2 (30.8%)	2C (28.2%)	1A4a (16.0%)	1A1a (8.2%)	1A4b (7.9%)	2B (3.4%)	6C (1.6%)						96.1
PAH	1A4b (38.7%)	2C (27.8%)	6C (20.1%)	1A4c (5.0%)	1A1c (4.0%)								95.5
Dioxin	1A2 (39.0%)	2C (27.8%)	1A4b (12.1%)	6C (11.5%)	1A4a (3.6%)	1A1a (2.1%)							96.2
HCB	6C (50.4%)	1A4a (17.6%)	1A3b i (17.4%)	1A3b ii (4.9%)	1A2 (3.2%)	1A3b iii (2.9%)							96.5
PCB	2C (47.8%)	1A1a (34.0%)	1A4a (10.7%)	1A2 (4.5%)									97.0

Color codes

1 Energy	3 Solvent and product use	6 Waste
2 Industry	4 Agriculture	7 Other

Table 1.4 Key categories for the Italian Emission Inventory in 2007

1.7 QA/QC and Verification methods

ISPRA has elaborated an inventory QA/QC procedures manual which describes specific QC procedures to be implemented during the inventory development process, facilitates the overall QA procedures to be conducted, as far as possible, on the entire inventory and establishes quality objectives (APAT, 2006). Specific QA/QC procedures and different verification activities implemented thoroughly the current inventory compilation are figured out in the annual QA/QC plans (ISPRA, 2009 [c]).

Quality control checks and quality assurance procedures together with some verification activities are applied both to the national inventory as a whole and at sectoral level. Future planned improvements are prepared for each sector by the relevant inventory compiler; each expert identifies areas for sectoral improvement based on his own knowledge and in response to different inventory review processes.

In addition to *routine* general checks, source specific quality control procedures are applied on a case by case basis, focusing on key categories and on categories where significant methodological and data revision have taken place or new sources.

Checklists are compiled annually by the inventory experts and collected by the QA/QC coordinator. These lists are also registered in the 'reference' database.

General QC procedures also include data and documentation gathering. Specifically, the inventory analyst for a source category maintains a complete and separate project archive for that source category; the archive includes all the materials needed to develop the inventory for that year and is kept in a transparent manner.

Quality assurance procedures regard different verification activities of the inventory.

Feedbacks for the Italian inventory derive from communication of data to different institutions and/or at local level. Emission figures are also subjected to a process of re-examination once the inventory, the inventory related publications and the national inventory reports are posted on website, specifically www.apat.gov.it.

The preparation of environmental reports where data are needed at different aggregation levels or refer to different contexts, such as environmental and economic accountings, is also a check for emission trends. At national level, for instance, emission time series are reported in the Environmental Data Yearbooks published by the Institute, in the Reports on the State of the Environment by the Ministry for the Environment, Land and Sea and, moreover, figures are communicated to the National Institute of Statistics to be published in the relevant Environmental Statistics Yearbooks as well as used in the framework of the EUROSTAT NAMEA Project.

Comparisons between national activity data and data from international databases are usually carried out in order to find out the main differences and an explanation to them. Emission intensity indicators among countries (e.g. emissions per capita, industrial emissions per unit of added value, road transport emissions per passenger car, emissions from power generation per kWh of electricity produced, emissions from dairy cows per tonne of milk produced) can also be useful to provide a preliminary check and verification of the order of magnitude of the emissions. Additional comparisons between emission estimates from industrial sectors and those published by the industry itself in the Environmental reports are carried out annually in order to assess the quality and the uncertainty of the estimates.

The quality of the inventory has also improved by the organization and participation in sector specific workshops.

A specific procedure undertaken for improving the inventory regards the establishment of national expert panels (in particular, in road transport, land use change and forestry and energy sectors) which involve, on a voluntary basis, different institutions, local agencies and industrial associations

cooperating for improving activity data and emission factors accuracy.

Furthermore, activities in the framework of the improvement of local inventories are carried out together with local authorities; the first meetings have already taken place concentrating on the comparison between top down and bottom up approaches identifying the main critical issues. In 2008, ISPRA finalised the provincial inventory at local scale for the years 1990, 1995, 2000 and 2005 (ISPRA, 2009 [d]) applying a top down approach. Methodologies and results were checked out by regional and local environmental agencies and authorities, and are also available at ISPRA web address <http://www.sinanet.apat.it/it/inventaria>. This work is also relevant to carry out regional scenarios, for the main pollutants, within the Rains Italy project implemented by ENEA and supported by ISPRA and the regional authorities.

In addition to these expert panels, ISPRA participates in technical working groups within the National Statistical System. These groups, named *Circoli di qualità*, coordinated by the National Institute of Statistics, are constituted by both producers and users of statistical information with the aim of improving and monitoring statistical information in specific sectors such as transport, industry, agriculture, forest and fishing. These activities should improve the quality and details of basic data, as well as enable a more organized and timely communication.

Other specific activities relating to improvements of the inventory and QA/QC practises in the last year regarded the progress on the building of a unique database where information collected in the framework of different European directives, Large Combustion Plant, E-PRTR and Emissions Trading, are gathered together thus highlighting the main discrepancies in information and detecting potential errors. Even though the database is not completed yet all the figures are considered in an overall approach and used in the compilation of the inventory.

A proper archiving and reporting of the documentation related to the inventory compilation process is also part of the national QA/QC programme.

All the material and documents used for the inventory preparation are stored at the Institute for Environmental Protection and Research.

Information relating to the planning, preparation, and management of inventory activities are documented and archived. The archive is organised so that any skilled analyst could obtain relevant data sources and spreadsheets, reproduce the inventory and review all decisions about assumptions and methodologies undertaken. A master documentation catalogue is generated for each inventory year and it is possible to track changes in data and methodologies over time. Specifically, the documentation includes:

- electronic copies of each of the draft and final inventory report, electronic copies of the draft and final NFR tables;
- electronic copies of all the final, linked source category spreadsheets for the inventory estimates (including all spreadsheets that feed the emission spreadsheets);
- results of the reviews and, in general, all documentation related to the corresponding inventory year submission.

After each reporting cycle, all database files, spreadsheets and electronic documents are archived as 'read-only' mode.

A 'reference' database is also compiled every year to increase the transparency of the inventory. This database consists of a number of records that references all documentation used during the inventory compilation, for each sector and submission year, the link to the electronically available documents and the place where they are stored as well as internal documentation on QA/QC procedures.

1.8 General uncertainty evaluation

An overall uncertainty analysis for the Italian inventory related to the pollutants described in this report has not been assessed yet. Nevertheless, different studies on uncertainty have been carried out (Romano et al., 2004) and a quantitative assessment of the Italian GHG inventory is performed by the Tier 1 method defined in the IPCC Good Practice Guidance (IPCC, 2000) which provides a calculation based on the error propagation equations. Details on the results of the GHG inventory uncertainty figures can be found in the National Inventory Report 2009 (ISPRA, 2009 [b]).

It should be noted that different levels of uncertainty pertain to different pollutants. Estimates of the main pollutants are generally of high level, but PM emissions, especially those of small particle sizes, heavy metal and POP estimates are more uncertain. For this reason, even though not quantified in terms of uncertainty, improvements are planned especially for the specified pollutants.

Nevertheless, since quantitative uncertainty assessments constitute a means to either provide the inventory users with a quantitative assessment of the inventory quality or to direct the inventory preparation team to priority areas, a planned improvement for next submissions is the completion of such analysis.

1.9 General Assessment of Completeness

The inventory covers all major sources, as well as all main pollutants, included in the EMEP CORINAIR guidelines.

NFR sheets are complete as far as the details of basic information are available.

Allocation of emissions is not consistent with the guidelines only where there are no sufficient data available to split the information. For instance, emissions from combustion in manufacturing industries and construction are not split among the relevant production sectors but included in the total category; emissions from category 1.A.5.a other, stationary are reported and included under category 1A4a commercial and institutional emission estimates. PAH emissions are not detailed in the four indicator compounds but accounted for as a total.

There are only a few emission sources not assessed yet: NO_x emissions from manure management, from cattle, buffalo, swine and other livestock categories, and NO_x emissions from direct soil emission, from the use of fertilizers in soils.

Other not estimated emissions are PCPs and SCCP from solvent use, deriving from wood preservation and some manufacturing industries, and pesticides in agriculture. No information on activity data and emission factors are available for these sources at the moment and verification is needed to assess if these emissions actually occur within the national area.

Further investigation will be carried out about these source categories and pollutants in order to calculate and improve figures.

2 ANALYSIS OF KEY TRENDS BY POLLUTANT

2.1 Main pollutants

In the following sections, Italian emission series of sulphur oxides, nitrogen oxides, non-methane volatile organic compounds, carbon monoxide and ammonia are presented.

2.1.1 Sulphur dioxide (SO_x)

The national atmospheric emissions of sulphur oxides have significantly decreased in recent years, as occurred in almost all countries of the UNECE.

Figure 2.1 and Table 2.1 show the emission trend from 1990 to 2007. Figure 2.1 also illustrates the share of SO_x emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.




Figure 2.1 SO_x emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
<i>Gg</i>										
Combustion in energy and transformation industries	1,001	776	467	415	373	283	258	187	184	143
Non industrial combustion plants	96	35	22	23	20	20	19	19	17	14
Combustion - Industry	299	215	104	95	91	87	77	73	65	65
Production processes	156	125	50	60	61	56	55	59	53	59
Road transport	131	72	12	13	11	12	12	2	2	2
Other mobile sources and machinery	100	86	84	82	50	50	50	51	48	46
Waste treatment and disposal	13	11	10	10	9	9	9	11	9	9
Total	1,795	1,319	749	697	616	518	480	401	379	338

Table 2.1 SO_x emission trend from 1990 to 2007 (Gg)

Figures show a general decline of SO_x emissions during the period, from 1,795 Gg in 1990 to 338 Gg in 2007. The national target of SO_x emissions amounts to 480 Gg for 2010, as set by the National Emission Ceilings Directive.

The decreasing trend is determined mainly by the reduction in emissions from *combustion in energy* (-86%) and *industrial sectors* (-78%), representing in 2007 about 42% and 19% of the total, respectively. Emissions deriving from *non industrial combustion plants* (-85%) and *road transport* (-99%) show a strong decrease too, but these emissions represent only about 4% and 1% of the total in 2007, respectively. *Production processes* and *other mobile sources and machinery* also present a significant decreasing trend, showing an influence on the total of 17% and 14% and dropping by about -62% and -54%, respectively.

Since SO_x emissions are included in the NEC directive, an explanation of the sectoral decreasing trend, starting from the early eighties, is outlined more in details in the following.

Combustion in energy and transformation industries

The trend of emissions of this sector shows a reduction in the early eighties mainly due to the use, in the energy production, of natural gas in place of coal, and to the implementation of the Directive EEC 75/716 which introduces more restrictive constraints in the sulphur content of liquid fuels.

During the years 1985-1990, there was an increase of energy consumption that, not sufficiently hampered by additional measures, led to an increase in the emissions of the sector and consequently of total SO_x levels.

However in the nineties, there was an inverse trend due to the introduction of two regulatory instruments: the DPR 203/88, laying down rules concerning the authorisation of plants, and the DM of 12th July 1990, which introduced plant emission limits. Also the European Directive 88/609/EEC

concerning the limitation of specific pollutants originated from large combustion plants (transposed in Italy by the DM of 8th May 1989), gave a contribution to the reduction of emissions in the sector. Finally, in recent years, a further shift to natural gas in place of fuel oil has contributed to a decrease in emissions.

Non industrial combustion plants

The declining of the emissions occurred mainly as a result of the increase in natural gas and LPG as fuel alternative to coal and fuel oil for heating; furthermore, a number of European Directives on the sulphur content in fuels were adopted. In accordance with national legislation, the sulphur content allowed in diesel fuel has decreased from 0.8% in 1980 to 0.2% in 1995, while in fuel oil for heating from 3% in 1980 to 0.3% in 1998.

Combustion in industry

Emissions from this sector show the same trend of reduction in the area previously analyzed, as both submitted to the same rules.

Production processes

Emissions from refineries have been reduced as a result of compliance with the DM 12th July 1990, which introduces limit values. The reduction of emissions from chemical industry is due to the drop off of the sulphuric acid production and to the decrease of emissions in the production of carbon black. Furthermore, there was a reduction in emissions in the production of cement with regard to the type of fuel used in the process and the respective sulphur content.

Road transport

The reduction of emissions is mainly due to the introduction of Directives regulating the sulphur content in liquid fuels.

Other mobile sources and machinery

As regards off roads, emissions mainly derive from maritime transport, which shows a decrease due the introduction of Directives regulating the sulphur content in fuels.

2.1.2 Nitrogen oxides (NO_x)

The national atmospheric emissions of nitrogen oxides show a decreasing trend in the period 1990-2007, from 2,003 Gg to 1,141 Gg. Figure 2.2 and Table 2.2 show the emission figures from 1990 to 2007. Figure 2.2 also illustrates the share of NO_x emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.




Figure 2.2 NO_x emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
<i>Gg</i>										
Combustion in energy and transformation industries	457	344	173	160	159	159	147	117	114	105
Non industrial combustion plants	62	64	75	79	78	86	91	95	92	90
Combustion - Industry	246	177	148	148	140	140	136	145	141	144
Production processes	30	31	9	9	12	13	14	16	13	11
Road transport	928	959	743	747	711	697	669	604	591	576
Other mobile sources and machinery	270	275	268	259	247	245	240	231	216	201
Waste treatment and disposal	9	14	13	14	13	14	17	15	14	14
Agriculture	0	1	0	0	0	0	0	0	0	0
Total	2,003	1,864	1,428	1,416	1,361	1,354	1,314	1,224	1,182	1,141

Table 2.2 NO_x emission trend from 1990 to 2007 (Gg)

Total emissions show a reduction of about 43% from 1990 to 2007, with a marked decrease between 1995 and 2000, especially in the road transport and energy combustion sectors. The target value of emissions, fixed for 2010 by the National Emission Ceilings Directive, amounts to 990 Gg. The main source of emissions is *road transport* (about 51% in 2007), which shows a reduction of 38% between 1990 and 2007; *other mobile sources and machinery* in 2007 contributes to the total emissions for 18% and have reduced by 26% from 1990. Combustion in energy and in industry shows a decrease of about 77% and 42%, respectively, having a share on the total of about 9% and 13%, respectively. Among the sectors concerned, the only ones which highlight an increase in emissions are: *waste treatment and disposal* and *non industrial combustion plants*, showing an increase by 56% and 44%, respectively, but accounting only for 1% and 8% of the total, respectively.

As SO_x, NO_x emissions are also included in the NEC directive. Details on the sectoral emission trend and respective variation are outlined in the following sections, starting from the early eighties.

Combustion in energy and transformation industries

Emissions from this sector show an upward trend until 1988 due to an increase in energy consumption, not prevented by reduction measures. From 1988 onwards, emissions present a gradual reduction due, mainly, to the introduction of the two regulatory instruments already mentioned for sulphur dioxide: the DPR 203/88, laying down rules for the authorization of facilities and the DM of 12th July 1990, which introduces plant emission limits. The adoption of these regulations, as the DM 8th May 1989 on large combustion plants, has led to a shift in energy

consumption from oil with high sulphur content to oil with lower sulphur content and to natural gas. In recent years, the conversion to the use of natural gas to replace fuel oil has intensified, thanks to incentives granted for the improvement of energy efficiency. These measures, together with those of promoting renewable energy and energy saving, have led to a further reduction of emissions in the sector.

Non industrial combustion plants

The increase in emissions is explained by the growing trend of energy consumption during the period considered. This is due the fact that from the last twenty years all the new buildings are equipped with heating system and old buildings were modernized.

Combustion in industry

Emissions from this sector show a decreasing trend, motivated by the same reasons as the energy industry, having undergone the same legislation.

Road transport

The decrease is the result of two opposing trends: an increase in emissions in the early years of the historical series, with a peak in 1992, due to the increase in the fleet and in the total mileage of both passengers and goods transported by road, and a subsequent reduction in emissions. This decrease is, once more, the result of two opposing trends: on the one hand, the growth of both the fleet and the mileage, on the other the introduction of technologies to reduce vehicle emissions, as the catalytic converter, provided by European Directives, in particular the Directives 91/441/EC, 94/12/EC and 98/69/EC on light vehicles.

To encourage the reduction of emissions, different policies have also been implemented, including incentives to renew the public and private fleet and for the purchase of electric vehicles, promotion for the integrated expansion of rail, maritime and urban transport system, and programmes of sustainable mobility.

Other mobile sources and machinery

From 1980, emissions have a slightly rising trend until 1998 and then decrease slightly until arriving in 2007 to lower levels. Emissions in the sector are characterized predominantly by maritime transport, by machinery used in agriculture and industry and to a lesser extent, by air transport. Regarding mobile machinery used in agriculture and industry, these sectors were not governed by any legislation until the Directive 97/68/EC, which provides for a reduction in NO_x limits from 1st January 1999, with a following decreasing trend particularly in recent years. Regarding aviation, in the absence of specific legislation up to now, emissions have increased in relation to the growth in air traffic.

2.1.3 Ammonia (NH₃)

The national atmospheric emissions of ammonia show a slight decline in the period 1990-2007, from 466 Gg to 418 Gg. Figure 2.3 and Table 2.3 report the emission figures from 1990 to 2007. Figure 2.3 also illustrates the share of NH₃ emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.

According to the National Emission Ceilings Directive, the target value of emissions for 2010 amounts to 420 Gg.




Figure 2.3 NH₃ emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
<i>Gg</i>										
Combustion in energy and transformation industries	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2
Non industrial combustion plants	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Combustion - Industry	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Production processes	0.8	0.4	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2
Road transport	0.8	6.9	24.7	23.8	22.5	20.3	19.1	17.7	16.7	14.2
Other mobile sources and machinery	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Waste treatment and disposal	7.0	8.3	9.1	9.2	8.9	8.6	8.2	8.2	7.9	7.9
Agriculture	457.5	432.5	416.7	418.5	407.4	404.2	398.6	387.5	384.7	395.5
Total	466.3	448.4	451.0	452.0	439.3	433.7	426.5	414.0	409.8	418.1

Table 2.3 NH₃ emission trend from 1990 to 2007 (Gg)

In 2007 *agriculture* is the main source of emissions, with a contribution by 95% out of the total NH₃ emissions; from 1990 to 2007 emissions from this sector decrease of about 14%. Emissions from *road transport* show a strong increase, but the share on the total is only about 3%. Emissions from *waste treatment and disposal*, accounting only for 2% of the total, increase of about 13%. Emissions from *combustion in energy and transformation industries* show an increase of about 41%, but in 2007 the contribution to total emissions is almost zero. Emissions from *non industrial combustion plants* decrease of about 81%, but the contribution to total emissions is negligible. Emissions from *combustion in industry* and *production processes* show a reduction of about 19% and 79% respectively, but also this contribution is insignificant.

Specifically, emissions from agriculture have decreased for both the merging of animal farms in large companies and the introduction of abatement technologies due to the implementation of the EU IPPC Directive. Emissions related to production processes, mainly the production of nitrogenous fertilizers and ammonia, dropped as a result of a lower production, whereas emissions from the waste sector have increased as a result of the greater amount of waste disposed in landfills. Emissions from road transport have increased as a result of the introduction of catalytic converter.

2.1.4 Non methane volatile organic compounds (NMVOC)

The national atmospheric emissions of NMVOC show a decreasing trend in the period 1990-2007. Figure 2.4 and Table 2.4 illustrate the emissions values from 1990 to 2007. Figure 2.4 also illustrates the share of NMVOC emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.




Figure 2.4 NMVOC emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
<i>Gg</i>										
Combustion in energy and transformation industries	7.6	7.4	6.3	5.6	5.7	5.6	5.5	5.6	5.6	5.6
Non industrial combustion plants	24.9	33.0	43.4	46.2	40.6	45.7	56.3	55.0	58.5	68.7
Combustion - Industry	7.3	8.1	8.2	7.9	7.7	7.8	8.0	8.0	8.3	8.2
Production processes	95.0	85.6	70.7	72.9	78.4	72.4	75.0	76.1	76.9	74.8
Extraction and distrib. of fossil fuels / geothermal energy	90.4	103.7	56.2	51.0	55.3	54.1	52.5	53.6	51.0	47.9
Solvent and other product use	589.9	545.1	512.3	505.2	504.4	500.0	501.9	499.9	512.5	515.0
Road transport	916.4	1,008.3	689.2	636.7	573.2	526.6	461.5	394.4	357.7	327.4
Other mobile sources and machinery	187.3	183.7	154.6	148.9	140.8	136.6	131.2	130.4	126.5	123.0
Waste treatment and disposal	19.0	25.7	23.9	25.1	24.6	24.3	27.0	25.3	24.3	23.5
Agriculture	1.3	1.3	1.2	1.2	1.2	1.2	1.3	1.2	1.2	1.2
Total	1,939.0	2,001.8	1,566.0	1,500.6	1,432.1	1,374.1	1,320.2	1,249.4	1,222.4	1,195.3

Table 2.4 NMVOC emission trend from 1990 to 2007 (Gg)

The global emission trend shows a reduction of about 38% between 1990 and 2007, from 1,939 Gg to 1,195 Gg. In the framework of the National Emission Ceilings Directive, the target value of NMVOC for 2010 has been fixed to 1,160 Gg.

Solvent and other product use is the main source of emissions, contributing to the total with 43% and showing a decrease of about 13%. The biggest reductions relate to the sector of *road transport* (-64%), accounting for 27% of the total and the sector of *extraction and distribution of fossil fuels/geothermal energy* (-47%), accounting only for 4%. Emissions from *non industrial combustion plants* show the biggest increase, but this is not relevant on total emissions, accounting only for 6%. Emissions from *waste treatment and disposal*, and *from combustion in industry*, accounting only for 2% and 1% of the total, show increase of about 24% and 12% respectively. Emissions from *other mobile sources and machinery*, accounting for 10% of the total, decrease of about 34%.

Details on the sectoral emission trend and respective variation are outlined in the following sections.

Solvent and other product use

Emissions from this sector stem from numerous activities such as painting both domestic and industrial, degreasing and dry cleaning, manufacturing and processing chemicals, other use of

solvents and related activities including the use of household products that contain solvents, such as cosmetics, household products and toiletries.

Significant reductions occurred in the nineties by the introduction in the market of products with low solvent content in paints, and the reduction of the total amount of organic solvent used for metal degreasing and in glues and adhesives; furthermore, there was a replacement of open loop with closed loop laundry machines. The gradual application of the EU Directive 99/13/EC will lead to further reductions in the coming years.

Road transport

The trend of emissions in this sector is characterized by a first stage of reduction in the early eighties, occurred despite the increase of consumption and mileage because of the gradual adjustment of the fleet to Community legislation, ECE Regulation 15 and subsequent amendments, which introduced stricter emission limits for passenger cars. Subsequently, in the early nineties, an increase in emissions is observed, with a peak in 1992, due to a high increase in gasoline consumption not efficiently opposed by the replacement of the fleet. With the introduction of Directive 91/441/EC and following, which provide for cars the catalytic device to reduce exhaust and evaporative emissions, NMVOC emissions were gradually reduced.

A different explanation of the emission trend pertains to the nineties. In fact, in this period an increase of the fleet and of the mileage is observed in Italy, especially for the emergent use of mopeds for urban mobility, which, until 1999, were not subject to any national emission regulation. Thereafter, various measures were introduced in order to facilitate the reduction of NMVOC emissions, including incentives for replacement of both the fleet of passenger cars and of mopeds and motorcycles with low-emission vehicles; incentives were also provided for the use of fuels different from gasoline, such as LPG and natural gas. In addition, funds were allocated for the implementation of urban traffic plans, for the establishment of restricted traffic areas and car-free days, for checks on exhaust pipes of cars, for the implementation of voluntary agreements with manufacturers of mopeds and motorcycles in order to anticipate the timing provided by the European Directive 97/24/EC as regards the placing on the market of mopeds with low emissions.

Other mobile sources and machinery

The reduction in emissions is explained by the reduction of gasoline consumption in the sector, largely for two-stroke engines used in agriculture and in maritime activities.

As regards the other sectors, a decrease in emissions from production processes is observed, mainly in the food industries, in the chemical sector and in the processes in the refineries. The emissions concerning the extraction and distribution of fuels, even in the presence of an increase in quantity treated, have been reduced as a result of the application of the DM 16th May 1996, concerning the adoption of devices for the recovery of vapours and of the applications of measures on deposits of gasoline provided by the DM 21st January 2000.

Emissions from the other sectors are not subject to specific regulations.

2.1.5 Carbon monoxide (CO)

The national CO emissions show a decreasing trend in the period 1990-2007, from 6,928 Gg to 3,336 Gg. The emission figures from 1990 to 2007 are shown in Figure 2.5 and Table 2.5. Figure 2.5 also illustrates the share of CO emissions by category in 1990 and 2007, as well as the total and sectoral variation from 1990 to 2007.




Figure 2.5 CO emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
<i>Gg</i>										
Combustion in energy and transformation industries	59	54	56	58	55	51	51	54	54	55
Non industrial combustion plants	256	348	450	469	395	428	506	479	524	658
Combustion - Industry	306	411	312	309	285	295	287	331	326	355
Production processes	215	124	114	110	106	113	117	121	128	119
Road transport	5,354	5,155	3,271	3,045	2,758	2,563	2,236	1,869	1,692	1,551
Other mobile sources and machinery	567	503	396	375	347	337	335	346	327	315
Waste treatment and disposal	159	269	249	271	263	269	339	296	281	269
Agriculture	13	13	12	11	12	11	14	13	12	13
Total	6,928	6,877	4,860	4,648	4,221	4,067	3,884	3,509	3,345	3,336

Table 2.5 CO emission trend from 1990 to 2007 (Gg)

The decrease in emissions (-52%) is mostly due to the trend observed for the transport sector (including road, railways, air and maritime transport) which show a global reduction from 1990 to 2007 of about 71%. Specifically by sector, emissions from *road transport* and *other mobile sources and machinery*, accounting in 2007 for 46% and 9% of the total, respectively, show a decrease from 1990 to 2007 of about 71% and 44%. On the other hand, emissions from *non industrial combustion plants*, representing about 20% of the total, show a strong increase between 1990 and 2007, equal to 157% due to the increase of wood combustion for heating; figures show a strong increase in emissions from *waste treatment and disposal* too (69%), which share 8% of the total.

2.2 Particulate matter

2.2.1 PM10

The national atmospheric emissions of PM₁₀ show a slight decreasing trend in the period 1990-2007, from 232 Gg to 162 Gg. Figure 2.6 and Table 2.6 illustrate the emission trend from 1990 to 2007. Figure 2.6 also illustrates the share of PM₁₀ emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.




Figure 2.6 PM₁₀ emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
Gg										
Combustion in energy and transformation industries	45	40	18	16	13	9	8	6	6	4
Non industrial combustion plants	14	16	20	21	17	19	22	20	22	28
Combustion - Industry	36	33	24	23	23	23	22	22	22	22
Production processes	22	20	19	19	18	19	20	20	20	20
Extraction and distrib. of fossil fuels / geothermal energy	1	1	1	1	1	1	1	1	1	1
Solvent and other product use	0	0	0	0	0	0	0	0	0	0
Road transport	57	56	52	52	50	48	46	42	40	39
Other mobile sources and machinery	31	33	29	28	27	27	25	24	21	19
Waste treatment and disposal	8	12	11	12	12	12	15	13	13	12
Agriculture	18	18	17	18	18	18	18	18	17	18
Total	232	229	192	191	179	175	177	165	162	162

Table 2.6 PM10 emission trend from 1990 to 2007 (Gg)

A considerable amount of emissions is mostly to be attributed to *road transport* (24% in 2007); from 1990 to 2007 the trend shows a reduction of about 32%. In 2007 *other mobile sources and machinery* account for 12% of the total and show a reduction of about 39%. Emissions from *non industrial combustion plants* and from *combustion in industry* account for about 17% and 14% of the total, respectively, but while the former show an increase of about 98%, the latter decrease of about 38%. Emissions from *production processes*, accounting for 12% of the total in 2007, decrease of about 10%, between 1990 and 2007. The largest decrease (-90%) is observed in emissions deriving from *combustion in energy and transformation industries*, which contribution to total emissions is equal to 3%.

2.2.2 PM2.5

The trend of the national atmospheric emissions of PM_{2.5} is slightly decreasing between 1990 and 2007, with a variation from 199 Gg to 130 Gg. Figure 2.7 and Table 2.7 illustrate the emission trend from 1990 to 2007. Figure 2.7 also illustrates the share of PM_{2.5} emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.




Figure 2.7 PM_{2.5} emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
<i>Gg</i>										
Combustion in energy and transformation industries	43	38	18	15	12	9	8	6	5	4
Non industrial combustion plants	12	14	19	19	16	18	21	19	21	26
Combustion - Industry	34	32	23	22	22	22	21	21	21	21
Production processes	11	10	8	8	8	8	8	8	9	9
Extraction and distrib. of fossil fuels / geothermal energy	1	1	1	1	1	1	1	1	1	1
Solvent and other product use	0	0	0	0	0	0	0	0	0	0
Road transport	54	52	48	48	45	44	42	37	36	35
Other mobile sources and machinery	31	33	29	28	27	27	25	24	21	19
Waste treatment and disposal	7	10	10	10	10	10	12	11	11	10
Agriculture	6	6	6	6	6	6	6	6	6	6
Total	199	195	161	158	147	143	144	133	130	130

Table 2.7 PM2.5 emission trend from 1990 to 2007 (Gg)

Total emissions show a global reduction from 1990 to 2007 of about 34%. Specifically, emissions from *road transport*, accounting for 27% of total emissions, decrease of about 36%. Emissions from *other mobile sources and machinery* show a reduction of 39%, accounting in 2007 for 15% of total emissions. Emissions from *non industrial combustion plants* and from *combustion in industry* account for 20% and 16% of the total, but while the former show an increase of about 116%, the latter decrease of about 38%. Emissions from *waste treatment and disposal*, accounting for 8% of the total in 2007, show an increase of about 46%. The largest decrease is observed for *combustion in energy and transformation industries* (-90%), being the influence on the total in 2007 equal to 3%.

2.3 Heavy metals (Pb, Cd, Hg)

This section provides an illustration of the most significant developments between 1990 and 2007 of lead, cadmium and mercury emissions.

2.3.1 Lead (Pb)

The national atmospheric emissions of lead show a strong decreasing trend (-94%) between 1990 and 2007, varying from 4,371 Mg to 273 Mg. Figure 2.8 and Table 2.8 illustrate the emission trend from 1990 to 2007. Figure 2.8 also illustrates the share of Pb emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.




Figure 2.8 Pb emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
<i>Mg</i>										
Combustion in energy and transformation industries	4	4	4	4	4	4	4	4	4	4
Non-industrial combustion plants	11	12	17	21	19	25	39	41	41	40
Combustion in manufacturing industry	263	235	153	149	139	137	135	142	142	143
Production processes	64	68	67	69	70	70	74	74	82	82
Road transport	3,880	1,617	678	447	0	0	0	0	0	0
Other mobile sources and machinery	144	46	13	9	1	1	1	1	1	1
Waste treatment and disposal	6	6	3	3	4	3	3	4	4	4
Total	4,371	1,988	935	702	236	241	256	265	273	273

Table 2.8 Pb emission trend from 1990 to 2007 (Mg)

In 2007 emissions from *processes with contact* have the most significant impact on the total (52%) and show a reduction of about 38%. Emissions from *production processes*, and in particular processes in iron and steel industries and collieries, increased of about 28%, and represent 30% of the total. Emissions from *non industrial combustion plants* show a strong increase and, in 2007, represent the 15% of the total. As regard emissions from *transport* activities, because of changes occurred in the legislation regarding fuels, trends show a sharp reduction in emissions from 2002 onwards. Emissions from *process furnaces without contact* show a strong decrease (-97%) but the contribution to total emissions in 2007 is negligible (equal to 0.4%).

2.3.2 Cadmium (Cd)

The national atmospheric emissions of cadmium show a slight decreasing trend. Figure 2.9 and Table 2.9 illustrate the emission trend from 1990 to 2007. Figure 2.9 also illustrates the share of Cd emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.




Figure 2.9 Cd emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
	<i>Mg</i>									
Combustion in energy and transformation industries	0.19	0.20	0.18	0.18	0.20	0.19	0.18	0.17	0.17	0.16
Non-industrial combustion plants	1.84	1.47	2.02	2.28	2.12	2.40	2.90	2.94	2.92	3.01
Combustion in manufacturing industry	5.61	5.56	4.98	4.55	3.00	3.01	3.07	3.28	3.36	3.28
Production processes	2.01	1.78	1.42	1.44	1.41	1.41	1.49	1.52	1.63	1.63
Road transport	0.30	0.32	0.34	0.35	0.35	0.36	0.37	0.36	0.37	0.37
Other mobile sources and machinery	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03
Waste treatment and disposal	0.31	0.27	0.14	0.16	0.18	0.14	0.13	0.16	0.17	0.17
Total	10.28	9.63	9.10	8.98	7.28	7.54	8.17	8.46	8.64	8.65

Table 2.9 Cd emission trend from 1990 to 2007 (Mg)

Emissions show a global reduction of 16% between 1990 and 2007, from 10.3 Mg to 8.7 Mg. Among the most significant variations, emissions from *combustion in industry* and from *non industrial combustion plants* represent the 38% and 35% of the total, showing the former a decrease (-42%) and the latter a strong increase (64%). Emissions from *production* decrease of about 19% and represent the 19% of the total. Emissions from *waste incineration*, accounting for 2% of the total, register a reduction of about 45%. The share of other subsectors on the total is irrelevant.

2.3.3 Mercury (Hg)

The national atmospheric emissions of mercury show a quite stable trend in the period 1990-2007. Figure 2.10 and Table 2.10 illustrate the emission trend from 1990 to 2007. Figure 2.10 also illustrates the share of Hg emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.




Figure 2.10 Hg emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
<i>Mg</i>										
Combustion in energy and transformation industries	1.10	1.15	1.10	1.12	1.21	1.18	1.15	1.10	1.09	1.06
Non-industrial combustion plants	0.90	1.08	1.50	1.71	1.57	1.86	2.42	2.48	2.52	2.65
Combustion in manufacturing industry	4.19	3.95	3.37	3.35	3.31	3.20	3.24	3.35	3.32	3.30
Production processes	5.31	4.26	3.50	3.47	3.34	3.13	3.40	3.31	3.59	3.52
Waste treatment and disposal	0.28	0.25	0.12	0.14	0.17	0.14	0.13	0.15	0.17	0.17
Total	11.79	10.70	9.59	9.79	9.59	9.51	10.34	10.39	10.70	10.71

Table 2.10 Hg emission trend from 1990 to 2007 (Mg)

Emission trend shows a global reduction of about 9% from 1990 to 2007, varying from 11.8 Mg to 10.7 Mg. The main variations concern: emissions from *processes in iron and steel industries and collieries*, representing 29% of the total and increasing of 27%; emissions from *processes with contact*, accounting for 26% and decreasing of 21%; emissions from *non industrial combustion plants* which represent 25% of the total and showing the strongest increase (193%). Emissions deriving from *combustion in energy and transformation industries*, accounting for 10%, show a reduction of 4%. Emissions from *processes in inorganic chemical industries*, contributing to the total only for 3%, show the largest reduction, equal to 87%.

2.4 Persistent organic pollutants (POPs)

In this section, the most significant peculiarities of polycyclic aromatic hydrocarbons and dioxins, occurred between 1990 and 2007, will be presented.

2.4.1 Polycyclic aromatic hydrocarbons (PAH)

The national atmospheric emissions of polycyclic aromatic hydrocarbons show an increasing trend between 1990 and 2007, from 104 Mg to 155 Mg. Figure 2.11 and Table 2.11 illustrate the emission trend from 1990 to 2007. Figure 2.11 also illustrates the share of PAH emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.




Figure 2.11 PAH emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
<i>Mg</i>										
Combustion in energy and transformation industries	9.09	7.68	6.54	6.93	5.97	5.62	5.82	6.41	6.56	6.52
Non-industrial combustion plants	23.11	32.72	44.56	46.38	39.59	42.93	51.81	48.68	53.78	68.68
Combustion in manufacturing industry	2.91	2.93	2.16	2.14	1.94	1.98	1.90	2.27	2.23	2.45
Production processes	44.48	44.06	42.44	39.90	37.27	38.88	40.93	43.81	44.51	43.16
Road transport	1.84	1.88	2.08	2.16	2.32	2.44	2.58	2.68	2.79	2.90
Other mobile sources and machinery	0.36	0.37	0.36	0.36	0.35	0.36	0.37	0.37	0.36	0.34
Waste treatment and disposal	21.28	31.10	29.54	31.22	30.34	30.36	37.81	33.90	32.50	31.23
Total	103.06	120.73	127.68	129.08	117.78	122.58	141.23	138.11	142.73	155.30

Table 2.11 PAH emission trend from 1990 to 2007 (Mg)

Between 1990 and 2007, global emissions show a growth of about 51%. Among the most significant changes, emissions from *residential plants* account for 39% of the total and show a strong increase (about 165%) due to the increase in wood consumption for heating; emissions from *processes in iron and steel industries and collieries* account for 28% of the total and show values quite stable between 1990 and 2007; emissions from *open burning of agricultural wastes* (except 10.03), accounting for 20% of the total, show an increase of 47%. Emissions from *plants in agriculture, forestry and aquaculture*, accounting for 5% in 2007, show a large growth from 2000 onwards, due to the use of biomass in plants. The share of other subsectors is less than 2%.

2.4.2 Dioxins

The national atmospheric emissions of dioxins show a decreasing trend between 1990 and 2007, with values varying from 472 g I Teq to 318 g I Teq. Figure 2.12 and Table 2.12 illustrate the emission trend from 1990 to 2007. Figure 2.12 also illustrates the share of dioxin emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.




Figure 2.12 Dioxin emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
<i>g I Teq</i>										
Combustion in energy and transformation industries	24.97	28.34	21.89	21.26	22.69	20.28	16.99	14.73	14.28	12.04
Non-industrial combustion plants	132.21	120.49	106.49	49.92	43.05	43.18	43.02	41.84	43.73	54.43
Combustion in manufacturing industry	117.29	121.13	110.65	108.40	103.03	103.87	102.87	116.25	115.70	124.07
Production processes	67.20	71.68	70.66	73.22	75.32	75.84	79.58	78.59	87.80	88.45
Road transport	7.22	7.45	4.19	3.90	3.49	3.31	2.88	2.57	2.48	2.48
Waste treatment and disposal	123.51	110.54	55.58	36.50	35.46	35.48	44.15	39.59	37.97	36.49
Total	472.40	459.61	369.46	293.20	283.03	281.96	289.49	293.57	301.96	317.96

Table 2.12 Dioxin emission trend from 1990 to 2007 (g I Teq)

The general trend shows a decrease from 1990 to 2007 equal to 33%, with a noticeable decline between 1995 and 2001. The most considerable reductions, between 1990 and 2007, are observed in *non-industrial combustion plants* and *waste treatment disposal* (-59% and -71%, respectively). Specifically, the reduction is principally due to the cut of emissions from the combustion of municipal waste both with energy recovery, reported under the non industrial sector, and without recovery, reported under the waste sector due to the introduction of regulations establishing more stringent limits of dioxin emissions from stacks.

In 2007, the subsector which has contributed most to total emissions is *combustion in manufacturing industries*, accounting for 39% of the total and showing an increase of 6% in the period 1990-2007. *Production processes* account for 28% of the total emissions in 2007 showing an increase of about 32% in the period 1990-2007.

2.4.3 Hexachlorobenzene (HCB)

The national atmospheric emissions of hexachlorobenzene show an increasing trend in the period 1990-2007, varying from 19 kg to 28 kg. Figure 2.13 and Table 2.13 illustrate the emission trend from 1990 to 2007. Figure 2.13 also illustrates the share of HCB emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.




Figure 2.13 HCB emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
<i>Mg</i>										
Combustion in energy and transformation industries	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Non-industrial combustion plants	0.001	0.002	0.006	0.014	0.014	0.014	0.005	0.005	0.006	0.005
Combustion in manufacturing industry	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
Road transport	0.003	0.003	0.004	0.004	0.005	0.005	0.006	0.006	0.007	0.007
Waste treatment and disposal	0.013	0.014	0.010	0.012	0.012	0.012	0.012	0.009	0.014	0.014
Total	0.019	0.020	0.021	0.030	0.032	0.033	0.024	0.021	0.028	0.028

Table 2.13 HCB emission trend from 1990 to 2007 (Mg)

The sector contributing most to the general trend is *waste incineration*, with exception of the years 2001-2003 where peaks are observed because of the relevant weight of the commercial sector due to the considerable increase of the amount of sludge incineration with energy recovery (which is accounted for in this sector) burnt in a specific incinerator. The other two relevant sectors are *road transport* and *non industrial combustion plants*, accounting for 25% and 19%, respectively; both sectors show a significant increase between 1990 and 2007.

2.4.4 Polychlorinated biphenyl (PCB)

The national atmospheric emissions of polychlorinated biphenyl show a slight decreasing trend in the period 1990-2007, about 1.8%, from 242 kg to 237 kg.

Figure 2.14 and Table 2.14 illustrate the emission trend from 1990 to 2007. Figure 2.14 also illustrates the share of PCB emissions by category in 1990 and 2007 as well as the total and sectoral variation from 1990 to 2007.




Figure 2.14 PCB emission trend, percentage share by sector and variation 1990-2007

	1990	1995	2000	2001	2002	2003	2004	2005	2006	2007
	<i>Mg</i>									
Combustion in energy and transformation industries	0.115	0.119	0.096	0.099	0.107	0.103	0.101	0.089	0.089	0.081
Non-industrial combustion plants	0.010	0.012	0.019	0.022	0.022	0.025	0.028	0.029	0.030	0.030
Combustion in manufacturing industry	0.019	0.016	0.012	0.012	0.012	0.013	0.012	0.011	0.011	0.011
Production processes	0.092	0.100	0.096	0.095	0.095	0.097	0.103	0.106	0.114	0.113
Waste treatment and disposal	0.006	0.005	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002
Total	0.242	0.252	0.226	0.231	0.237	0.240	0.245	0.237	0.246	0.237

Table 2.14 PCB emission trend from 1990 to 2007 (Mg)

The subsectors contributing most to the general trend are the *production processes* sector and the *combustion in energy and transformation industries* sector, accounting for 48% and 34% of the total emissions, respectively, and showing a reduction by 44% and 30%. The other relevant subsectors are *non industrial combustion plants* and *manufacturing industries and construction*, accounting for 12% and 4%, respectively; both sectors show an increase between 1990 and 2007.

3 ANALYSIS OF KEY TRENDS BY SECTOR

3.1 Energy (NFR SECTOR 1)

3.1.1 Methodological issues

Methodologies used for estimating emissions from this sector are based on and conform to the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007), the IPCC Guidelines (IPCC, 1997; IPCC, 2006) and the Good Practice Guidance (IPCC, 2000).

Specifically, for road transport, COPERT 4 programme has been used to calculate emissions (EEA, 2007); the updated version of the model has been applied for the whole time series, resulting, in particular, in an increase of NO_x emission levels.

A detailed description on the methods and national specific circumstances as well as reference material is documented in the national inventory report of the Italian greenhouse gas inventory (ISPRA, 2009[b]). At national level, trends of the CLRTAP pollutants are described in the environmental data yearbook published by ISPRA (ISPRA, 2009 [e]).

The National Energy Balance, published by the Ministry of Economic Development, is the main source of information to estimate emissions from the energy sector as it reports fuel consumption for different sectors at national levels. Additional information for electricity production is provided by the major national electricity producers and by the major national industry corporation. On the other hand, basic activity data for road transport, maritime and aviation, such as the number of vehicles, harbour statistics and aircraft landing and take-off cycles are provided in statistical yearbooks published both by the National Institute of Statistics and the Ministry of Transportation. Other data are communicated by different category associations.

The Institute, specifically the same unit responsible for the inventory compilation, also collects data in the framework of the European Emissions Trading Scheme, the National Pollutant Emission Register (EPER, now E-PRTR) and the Large Combustion Plants (LCP) Directives.

A unique database is being completed for the analysis of this information, so that figures are cross checked and used to develop country-specific emission factors and input activity data levels. These processes have improved the efficiency in collecting data and exchange of information, and whenever data cannot be straight used for the inventory compilation, they are taken into account as verification.

A complete description of methodological and activity data improvements are documented every year in a QA/QC plan (ISPRA, 2009[c]).

The following sections present an outline of the main key categories in the energy sector. Table 3.1 highlights the key categories identified in the sector.

	1A1a	1A1b	1A1c	1A2	1A3a ii (i)	1A3a i (i)	1A3b i	1A3b ii	1A3b iii	1A3b iv	1A3b v	1A3b vi	1A3c	1A3d ii	1A3e	1A4a	1A4b	1A4c	1A5b	1B1a	1B1b	1B2	
	%																						
SO_x	24.10	14.26	4.01	19.34	0.05	0.07	0.32	0.10	0.13	0.01			0.00	13.36	0.00	1.62	2.50	0.04	0.04			12.04	
NO_x	5.96	2.16	0.93	14.80	0.16	0.21	21.00	6.93	21.89	0.68			0.36	7.50	0.14	3.45	4.25	6.49	0.87			0.44	
NH₃	0.05	0.00	0.00	0.01			3.26	0.10	0.03	0.02			0.00	0.00		0.00	0.00	0.00	0.00				
NMVOC	0.35	0.08	0.05	1.03	0.05	0.10	6.75	0.81	1.41	16.39	2.04		0.04	8.10	0.00	1.27	4.14	1.78	0.20	0.20	6.23		
CO	0.94	0.13	0.57	11.09	0.07	0.11	23.26	2.33	1.70	19.20			0.03	5.84	0.01	0.55	17.19	3.71	1.24		0.03		
PM10	1.45	0.64	0.54	14.83	0.01		6.92	4.68	4.57	1.89		6.04	0.30	3.95	0.02	0.72	14.49	7.13	0.76	0.49	0.15	0.38	
PM2.5	1.71	0.76	0.64	17.63	0.01		8.62	5.83	5.69	2.35		4.11	0.37	4.89	0.02	0.86	17.12	8.88	0.94	0.61	0.15	0.47	
Pb	1.17	0.19	0.01	52.19	0.14	0.22	0.00	0.00	0.00	0.00			0.00	0.05		14.25	0.30	0.02	0.00		0.38		
Cd	1.47	0.35	0.02	37.99	0.00	0.00	2.62	0.66	0.83	0.16			0.00	0.20		24.17	9.55	1.10	0.00		2.73		
Hg	8.22	1.52	0.18	30.81												16.01	7.93	0.84			1.32		
PAH	0.22	0.02	3.96	1.62	0.00	0.00	1.18	0.29	0.34	0.06			0.01	0.05		0.70	38.67	4.97	0.01				
Dioxin	2.14	1.65	0.00	39.02			0.28	0.04	0.13	0.33						3.64	12.08	1.40					
HCB	1.74			3.24			17.42	4.93	2.89	0.08						17.64	1.51						
PCB	34.02	0.06	0.00	4.51												10.71	1.81	0.22					

Note: grey shaded are key categories

Table 3.1 Key categories in the energy sector in 2007

The *energy* sector is the main source of emissions in Italy with a share of more than 80% in different pollutants under the UNECE convention; specifically, for the main pollutants, in 2007 the sector accounts for:

- 98% in national total NO_x emissions;
- 88% in national total CO emissions;
- 92% in national total SO_x emissions.

Moreover, the sector comprises 82% of total PM_{2.5} emissions and is also an important source for heavy metals; specifically in 2007 energy sector is responsible for 82% of total Cd emissions and accounts for a high share of other heavy metals, i.e As (99%), Cu (93%), Ni (95%), Se (92%).

There are no particular differences as compared the sectoral share in 1990, except for lead whose contribution in 1990, accounting for 98% of total emissions, was about 30% higher than 2007.

The most important source of emissions in the sector, in 2007, is represented by *road transport* at least for the main pollutants: NO_x (51.0%), CO (46.5%), NMVOC (25.4%), particulate matter (PM₁₀ 18.0%, PM_{2.5} 22.5%) and HCB (25.3%). There has been a strong reduction of lead emissions from 1990 to 2007 in road transport due to replacement of lead gasoline.

Manufacturing industries and *construction* is the main source for heavy metals, accounting for more than 52% of lead total emissions, 38% for cadmium and 31% for mercury, and dioxin (39%). The source is also relevant for PM₁₀ and PM_{2.5}, as well as SO_x and NO_x, about 15-20% of total emissions.

Public electricity and *heat production* is the main source of SO_x emissions in 2007 with a share of 24.1%, followed by *manufacturing industries and construction*, as already reported, *petroleum refining* (14.3%) and national *navigation* (13.4%); the source is also important for PCB emissions (34%).

A sector which seems of increasing importance is the *non-industrial combustion*, as for NO_x, CO, PM, emissions, accounting for 15-25%, and PAH (44%); these emissions are prevalently due to *biomass combustion* which occurs in the winter period becoming critical for air quality issues. This source is a key category for heavy metals, HCB and PCB due to the increase of combustion of waste with energy recovery reported under the sector.

3.2 Industrial Processes (NFR SECTOR 2)

3.2.1 Methodological issues

Methodologies used for estimating emissions from this sector are based on and conform to the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007), the IPCC Guidelines (IPCC, 1997; IPCC, 2006) and the Good Practice Guidance (IPCC, 2000).

Moreover, as for the *energy* sector, a lot of information derives from data collected in the framework of the EPER/E-PRTR registry, Large Combustion Plant directives and European Emissions Trading Scheme. Other small plants communicate their emissions which are also considered individually. These processes have improved the efficiency in collecting data and exchange of information, and whenever data cannot be straight used for the inventory compilation, they are taken into account as verification practice. Environmental Reports published by industrial associations are also considered in the verification process.

A detailed description on the methods and national specific circumstances as well as reference material is documented in the national inventory report of the Italian greenhouse gas inventory (ISPRA, 2009[b]). At national level, trends of the CLRTAP pollutants are described in the environmental data yearbook published by ISPRA (ISPRA, 2009[e]). A complete description of methodological and activity data improvements are documented every year in a QA/QC plan (ISPRA, 2009[c]).

The following sections present an outline of the main key categories in the *industrial processes* sector. Table 3.2 reports the key categories identified in the sector.

	2A1	2A2	2A5	2A6	2B1	2B2	2B3	2B5	2C	2D1	2D2	2G
	%											
SO_x	4.19				0.00			2.44	1.23	0.07	0.07	
NO_x					0.05	0.04	0.00	0.19	0.28	0.01		
NH₃					0.00	0.00		0.03				
NMVOC			0.00	0.80	0.01			0.30	0.32	0.15	2.28	
CO					0.00			0.33	3.21			
PM10	3.78	1.16	0.06	1.63				0.45	4.58	0.02	0.01	
PM2.5	0.71	0.22	0.01	0.30				0.27	4.51	0.00	0.00	
Pb									28.56			0.8
Cd								1.05	15.10			
Hg								3.38	28.19			
PAH									27.79			
Dioxin									27.82			
HCB												
PCB									47.78			

Note: grey shaded are key sources

Table 3.2 Key categories in the industrial processes sector in 2007

There is a general reduction of emissions in the period for most of the pollutants due to the implementation of different directives at European and national level. A strong decrease is observed especially in the *chemical industry* due to the introduction of relevant technological improvements; this sector is still a key source for Hg emissions from chlorine production.

The most important source of emissions in the sector, in 2007, is represented by *metal production*, specifically iron and steel, at least for particulate matter, heavy metals and POPs.

For SO_x and particulate matter, significant emissions derive from *cement production*.

3.3 Solvent and other product use (NFR SECTOR 3)

3.3.1 Methodological issues

Methodologies used for estimating emissions from this sector are based on and conform to the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007), the IPCC Guidelines (IPCC, 1997; IPCC, 2006) and Good Practice Guidance (IPCC, 2000).

The sector is characterized by a multitude of activities which implies that the collection of activity data and emission factors is laborious. A lot of contacts have been established in different sectors with industrial associations and documentation collected even though improvements are still needed especially in some areas.

Country specific emission factors provided by several accredited sources are used, together with some data provided by the national EPER/E-PRTR Registry. Specific surveys based on local and regional inventories have been funded by ISPRA to check NMVOC emission factors and update emission estimates.

A detailed description on the methods and national specific circumstances, as well as reference material, is documented in the national inventory report of the Italian greenhouse gas inventory (ISPRA, 2009[b]). At national level, trends of the CLRTAP pollutants are described in the environmental data yearbook published by ISPRA (ISPRA, 2009[e]).

	3A	3B	3C	3D
	%			
SO _x				
NO _x				
NH ₃				
NMVOOC	18.75	1.83	6.57	15.93
CO				
PM10			0.01	
PM2.5			0.01	
Pb				
Cd				
Hg				
PAH				0.01
Dioxin				
HCB				
PCB				

Note: grey shaded are key sources

Table 3.3 Key categories in the solvent and other product use sector in 2007

The general reduction observed in the emission trend of the sector is due to the implementation of the European Directive 1999/13/EC on the limitation of emissions of volatile organic compounds due to the use of organic solvents, entered into force in Italy in January 2004, which establishes a reduction of the solvent content in products.

The main source of emissions, for NMVOC, is *paint application* where emissions derive mainly from construction and building and wood application. The second source of emissions, *other paint*, is for the most part characterized by emissions deriving from domestic solvent use.

3.4 Agriculture (NFR SECTOR 4)

3.4.1 Methodological issues

Methodologies used for estimating national emissions from this sector are based on and conform to the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007), the IPCC Guidelines (IPCC, 1997; IPCC, 2006) and the Good Practice Guidance (IPCC, 2000). Therefore, consistency among methodologies for the preparation of the agriculture emission inventory under the CLRTAP and UNFCCC is guaranteed and synergies among international conventions/European directives are implemented (Córdoba and De Laurentis, 2007).

The procedure for the estimation of NH₃ emissions has been initially reported in APAT (APAT, 2005). However, a detailed and updated description of the methodologies for the estimation of NH₃ emissions, as well as of national specific circumstances and reference material, is reported in the National Inventory Report for greenhouse gases (ISPRA, 2009[b]). At national level, trends of the CLRTAP pollutants are described in the environmental data yearbook published by ISPRA (ISPRA, 2009[e]) and in Córdoba *et al.*, 2008 for NH₃ emissions. Moreover, every 5 years the national agriculture UNFCCC/CLRTAP emission inventory is disaggregated at NUTS 3 level as requested by CLRTAP (Córdoba *et al.*, 2008). A database with the time series for all sectors and pollutants has been published (ISPRA, 2008[a]; ISPRA, 2008[b]; ISPRA, 2009[d]). Furthermore, a complete description of methods and activity data improvements are documented every year in a QA/QC plan (ISPRA, 2009[c]).

Since the 2006 submission, results from a specific project on Mediterranean area, the *MeditAIRaneo* project, have been included in the preparation of the CLRTAP/UNFCCC agriculture emission inventory (CRPA, 2006[a]). Moreover, outcomes from the convention signed between APAT and the Ministry for the Environment, Land and Sea on NH₃ emission scenarios have been incorporated (CRPA, 2006[b]; ENEA, 2006).

In the future, the implementation of an *ad hoc* “Survey on Agricultural Production Methods” (SAPM), regulated by the European Commission, will be crucial for improving the preparation of the national agriculture inventory. This survey will be carried out in Italy during the 2010 General Agricultural Census. Information such as animal grazing period, animal housing and storage systems characteristics, and use of manure/slurry for land application will be collected. Some information at provincial level (NUTS3) has been already collected with the incorporation of specific queries in the Farm Structure Survey (FSS) from 2005 and 2007.

The following sections present an outline of the main key categories in the agriculture sector. Table 3.4 reports the key categories identified in the agriculture sector.

	4B1a	4B1b	4B2	4B3	4B4	4B5	4B6	4B7	4B8	4B9	4B13	4C	4D1	4F	4G
	%														
SO_x															
NO_x															
NH₃	15.75	16.98	2.05	0.43	0.05		0.24	0.03	8.90	7.44	2.66		40.07		
NMVOC	0.01	0.02	0.00	0.00	0.00		0.00	0.00	0.01						0.05
CO															
PM10	0.68	0.84	0.07				0.05	0.01	2.23	5.73					1.35
PM2.5	0.54	0.69	0.06				0.04	0.00	0.46	0.93					1.68
Pb															
Cd															
Hg															
PAH															
Dioxin															
HCB															
PCB															

Note: grey shaded are key sources

Table 3.4 Key categories in the agriculture sector in 2007

The agriculture sector is the main source of NH₃ emissions in Italy with a share of more than 94%; for the main pollutants, in 2007 the sector accounts for:

- 94% in national total NH₃ emissions;
- 11% in national total PM₁₀ emissions;
- 4% in national total PM_{2.5} emissions.

Moreover, the sector comprises 0.4% of total CO emissions, 0.1% of NMVOC and 0.04% of NO_x. There are no particular differences as compared the sectoral share in 1990.

Concerning NH₃ emissions, the category *Manure Management (4B)* represent, in 2007, 54% of ammonia emissions (58% in 1990). In particular, NH₃ emissions from *cattle (4B1)* stand for 60% of the category emissions, while emissions from *swine (4B8)* and *poultry (4B9)* represent 16% and 14%, respectively. *Direct soil emissions (4D)*, especially for the use of chemical fertilizers, represent 40% in 2007 of ammonia emissions (41% in 1990).

Regarding PM₁₀ emissions, the category *Manure Management (4B)* accounts for 9.6% in 2007 (6.8% in 1990). *Poultry (4B9)* and *swine (4B8)* represent the major contributors to the total PM₁₀ emissions from category 4B (60% and 23%, respectively). The presence of large poultry and swine farms in the Po river basin assume a particular relevance, at regional level, for air quality issues especially for the specific meteorological conditions for that regional area.

Similar consideration may be done for PM_{2.5} emissions; the category *Manure Management (4B)* contributes for 2.7% in 2007 (1.9% in 1990). *Cattle (4B1)* accounts for 47%, while *poultry (4B9)* stands for 32% to the total PM_{2.5} emissions from category 4B.

3.5 Waste (NFR SECTOR 6)

3.5.1 Methodological issues

Methodologies used for estimating emissions from this sector are based on and conform to the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007), the IPCC Guidelines (IPCC, 1997; IPCC, 2006) and the Good Practice Guidance (IPCC, 2000).

Activity data are derived by information from the waste cadastre, which is formed by a national branch hosted by ISPRA and regional and provincial divisions, and other national statistics. A complete database of the national incineration plants is available and updated annually, which reports the year of the construction and possible upgrade, the typology of combustion chamber and gas treatment section, if it is provided of energy recovery (thermal or electric), and the type and amount of waste incinerated (municipal, industrial, etc.).

Emission parameters and emission factors are derived by national studies or default figures, according to data availability.

A detailed description on the methods and national specific circumstances as well as reference material is documented in the National Inventory Report on the Italian greenhouse gas inventory (ISPRA, 2009 [b]).

The following sections present an outline of the main key categories in the waste sector.

	6A	6B	6C	6D
	%			
SO_x			0.08	
NO_x			1.17	
NH₃	1.62			0.26
NMVOC	0.70		1.21	0.03
CO			8.07	
PM10			7.40	
PM2.5			7.91	
Pb			1.52	
Cd			2.00	
Hg			1.60	
PAH			20.11	
Dioxin			11.48	
HCB			50.36	
PCB			0.89	

Note: grey shaded are key sources

Table 3.5 Key categories in the waste sector in 2007

The waste sector, and in particular *Waste incineration* (6C), is a relevant source of different pollutants; for the main pollutants, in 2007, the sector accounts for:

- 50% in national total HCB emissions;
- 20% in national total PAH emissions.
- 11% in national total Dioxin emissions;

Moreover, the sector comprises 7.4% and 7.9% of total PM10 and PM2.5 emissions, respectively, 8.1% of CO, 1.2% of NO_x, and more than 1.5% of heavy metals (HM). Comparing the sectoral 1990 emissions, it's possible to note a severe reduction of the share of HCB (69.4% in 1990), and a decrease in the share of dioxin emissions (26.1% in 1990) as a consequence of the introduction of more stringent limits of these emissions for incineration plants.

The European Council Directive 99/31/EC on the landfill of waste transposed by the Legislative Decree 13 January 2003 n.36, has been applied to the Italian landfills since July 2005, but the effectiveness of the policies will be significant in the future.

4 Recalculations and Improvements

4.1 Recalculations

To meet the requirements of transparency, consistency, comparability, completeness and accuracy of the inventory, the entire time series from 1990 onwards is checked and revised every year during the annual compilation of the inventory. Measures to guarantee and improve these qualifications are undertaken and recalculations should be considered as a contribution to the overall improvement of the inventory.

Recalculations are elaborated on account of changes in the methodologies used to carry out emission estimates, changes due to different allocation of emissions as compared to previous submissions, changes due to error corrections and in consideration of new available information.

The complete NFR files from 1980 to 2007 have been submitted.

The percentage difference between the time series reported in the 2008 submission and the series reported this year (2009 submission) are shown in Table 4.1 by pollutant.

Improvements in the calculation of emission estimates have led to a recalculation of the entire time series of the national inventory. Considering the total emissions, the emission levels for the year 2006 showed a decrease especially for CO, PM₁₀, PM_{2.5} and SO_x.

Relevant changes in the whole time series regarded, in particular, a revision of the emission estimates from the transport sector. Specifically, for road transport, the new version of Copert model, Copert 4 (EEA, 2007), has been applied to calculate emissions of all pollutants for the whole period 1990-2007. The recalculation affected mainly NO_x figures, which show an increase of about 4% of national total with respect to the previous submission. Other minor recalculation due to the application of Copert 4 concerned CO, NMVOC, NH₃, PM and Cd emission levels. For the maritime sector, emissions have been revised from 1998 on account of the results of a national study (TECHNE, 2009), which considered most recent trends in maritime activities in terms of technological improvements, fleet composition and relevant fuel consumption, and operational times especially hotelling and manoeuvring in harbour activities; the use of new information led to a reduction of ship average fuel consumptions for the last years. Besides it led to a reduction of CO and NMVOC emissions and an increase of NO_x emissions in harbour for the whole time series. Finally, changes also affected emissions from the aviation sector, from 2000, because of a revision of consumption values and relative parameters of the aviation fleet on the basis of the results of the national study (TECHNE, 2009), which considered most recent technological improvements in civil aviation, fleet composition and a more accurate split between national and international fuel consumption. It resulted in a general decrease of average fuel consumption and CO, NMVOC, PM and NO_x emissions.

As for the other sectors and pollutants, a revision of coke and sinter plants emission factors lead to a recalculation of PAH emissions, which levels increased from about 10% in 1990 to 6% in 2006. The high revision of HCB emissions in 2006 is due to the update of activity data for incineration plants which treat industrial wastes in addition to the above mentioned application of Copert 4 for road transport emissions.

Minor changes affected NH₃ emissions, for the whole time series, due to revision of emission factors for cattle and buffalo for the spreading phase.

	SO _x	NO _x	NH ₃	NM VOC	CO	PM10	PM2.5	Pb	Hg	Cd	DIOX	PAH
	%											
1980	0.0	1.8	0.4	-0.1	-1.9							
1981	0.0	1.8	0.4	-0.1	-1.9							
1982	0.0	1.9	0.4	-0.1	-1.9							
1983	0.0	1.9	0.4	-0.1	-1.9							
1984	0.0	1.9	0.4	-0.1	-1.9							
1985	0.0	1.8	0.4	0.0	-2.0							
1986	0.0	3.4	0.4	-1.4	-2.8							
1987	0.0	3.2	0.4	-0.8	-2.8							
1988	0.0	3.1	0.4	-1.1	-2.8							
1989	0.0	3.2	0.4	-0.9	-3.0							
1990	0.0	3.1	0.4	-2.1	-2.8	-5.8	-5.5	-0.2	0.0	2.2	0.0	10.6
1991	0.1	2.9	0.4	-2.1	-3.4	-6.3	-6.1	0.4	0.0	2.0	-0.1	9.0
1992	0.1	2.9	0.3	-3.2	-3.5	-6.8	-6.5	0.8	0.0	2.3	-0.1	8.4
1993	0.0	3.2	0.3	-0.5	-2.0	-5.9	-5.6	1.4	0.0	2.4	0.0	7.9
1994	0.0	3.0	0.3	0.5	-3.0	-5.6	-5.2	2.2	0.0	2.5	0.0	8.0
1995	0.0	3.0	0.4	-0.1	-4.0	-5.2	-4.7	3.0	0.0	2.6	0.0	7.9
1996	0.0	3.8	0.4	0.6	-4.3	-5.4	-4.9	3.7	0.0	2.6	0.0	8.0
1997	0.0	4.1	0.5	1.3	-4.8	-5.3	-4.7	4.8	0.0	2.7	0.0	7.6
1998	-0.2	4.3	0.5	2.9	-4.6	-4.6	-4.0	6.4	0.0	2.9	0.1	7.7
1999	-0.4	4.5	0.8	2.5	-6.9	-4.2	-3.4	8.4	0.0	3.0	0.0	6.9
2000	-0.8	3.8	2.2	4.5	-5.4	-4.0	-3.1	0.0	0.0	2.9	0.0	6.2
2001	-1.1	4.6	1.7	5.0	-8.8	-4.4	-3.6	0.0	-0.2	2.7	0.0	6.5
2002	-1.0	7.6	1.2	7.1	-5.3	-4.7	-3.8	-0.2	0.0	3.8	0.1	6.1
2003	-1.5	7.7	1.0	6.2	-6.9	-5.1	-4.3	-0.3	0.0	3.7	0.0	5.6
2004	-1.8	10.2	0.9	4.7	-7.7	-4.6	-3.7	-0.2	0.0	3.4	0.0	5.1
2005	-1.7	9.2	0.6	3.0	-8.5	-5.3	-4.4	-0.2	0.1	3.4	0.0	5.9
2006	-2.6	10.2	0.3	4.0	-6.9	-5.8	-4.9	-0.5	-0.1	3.1	0.0	5.9

Table 4.1 Recalculation between 2008 and 2009 submissions

4.2 Planned improvements

Specific improvements are specified in the 2009 QA/QC plan (ISPRA, 2009 [c]); they can be summarized as follows.

For the *energy* and *industrial processes* sectors, a major progress will regard the completion of a unique database where information collected in the framework of different directives, Large Combustion Plant, E-PRTR and Emissions Trading, are gathered together thus highlighting the main discrepancies in information and detecting potential errors. For the *agriculture* and *waste* sectors, improvements will be related to the availability of new information on emission factors, activity data as well as parameters necessary to carry out the estimates; specifically, a study on the best available technologies used in agriculture practises and availability of information on waste composition and other parameters following the entering into force of the European landfill directive.

Finally, efforts will be addressed to the comparison between local inventories and national inventory.

Further analyses will concern the collection of statistical data and information to estimate uncertainty in specific sectors.

5 Projections

The national projections reported within the UNECE Convention are calculated by the model RAINS Italy, the Italian version of the RAINS Europe model (Amman et al., 1999; IIASA, 2008). The estimations of SO₂, NO_x, NMVOC and NH₃ are based on an assessment of economic activities and a control strategy, explained by economic sector, set of abatement technologies planned in terms of rates of application for the current and future years (Pignatelli et al., 2007). Emission factors are those used for the national emission inventory estimations as well as national references and personal communication with sectoral experts.

In order to assess future economic activities levels two scenarios are developed:

- an energy scenario to estimate emissions from energy sources. The Markal (MARKet Allocation) model (Goldstein et al., 1999) is used to implement the scenario at 2010. Actually, this model has been modified at the beginning of the 1990s to take into consideration the Italian circumstances and evaluate potential and costs of emissions reduction of CO₂, NO_x e SO_x. Markal Italy (Gracceva and Contaldi, 2004) is also used to develop the energy mitigation scenario also for the Fourth National Communication under the UN Convention on Climate Change (MATTM, 2007).
- a scenario on production activities to estimate emissions from non energy sources. National statistics and projections of non energy economic activities are used to this end.

In addition to these scenarios, the national control strategy, that's the whole set of abatement technological measures to be implemented in the time interval considered, need to be defined.

Other documentation on emission scenarios in Italy can be found in Vialetto et al. (2005), Zanini et al. (2005).

Emission projections at 2010 for the pollutants regulated by the National Emission Ceilings (NEC) Directive (2001/81/EC) are reported in Table 5.1.

Moreover in Table 5.2, the national emission ceilings for Italy established by the D.Lgs. 171/2004 according to Annex I of the NEC directive are reported.

NFR2 codes	NO_x	SO_x	NH₃	NMVOC
			<i>kt</i>	
1A1a: Public electricity and heat	124.03	57.73	3.18	6.96
1A1b,c: Other energy industries	31.70	72.16	0.09	1.13
1A2: Manufacturing industries & construction	88.53	46.08	0.21	6.10
1A2a: Iron and steel	5.07	8.87		
1A2f: Other manufacturing industry	49.53	23.73		
1A3a: Civil aviation (LTO)	7.25	31.29		1.21
1A3b: Road transportation	424.97	0.56	5.87	225.73
1A3c: Railways	2.44	0.71	0.00	0.33
1A3d: Navigation	125.17	66.94	0.05	8.51
1A4a: Commercial / Institutional	73.56	10.61	1.77	
1A4b: Residential	0.29	2.35	0.01	113.32
1A4c: Agriculture & forestry	107.39	38.81	0.06	47.97
1B2a: Oil	8.63	1.11		44.64
1B2b: Natural gas				30.31
1B2c: Flaring in oil and gas extraction	1.11	8.86	0.00	
2B: Chemical industry	1.50	3.26	0.89	4.46
2D: Other production				34.69
3A: Paint application				155.71
3B: Degreasing and dry cleaning				2.92
3C: Chemical products, manufacturing and processing				33.76
3D: Other				201.56
4B1: Cattle			184.41	
4B3-7 & 13: Other			18.71	
4B8: Swine			47.97	
4B9: Poultry			51.74	
4D1: Direct soil emissions			89.61	
4F: Field burning of agricultural waste	4.97	2.18		12.34
6A: Solid waste disposal on land				9.34
6B: Waste water handling			4.95	
6C: Waste incineration	0.66	0.29		
7: Other			6.93	
Total	1,056.77	375.50	416.45	940.97

Table 5.1 Emission projections for the year 2010

NO _x	SO _x	NH ₃	NM VOC
		kt	
990	475	419	1159

Table 5.2 National emission ceilings for Italy

The latest projections show that the 2010 emissions ceilings will be reached for all the pollutants except for NO_x. Nevertheless, if the same methodology, as that for the 1998 negotiated for the NEC directive, is used NO_x projections would be 865 kt, whereas, taking into account the successive modifications in the EMEP CORINAIR methodology, figures reach 1,057 kt.

In 2009, a new energy scenario will be finalised and emission projections will be updated jointly with the preparation of projections of GHG emissions in the framework of the preparation of the fifth National Communication to the UNFCCC and the EU GHG Monitoring Mechanism requirements.

An accurate comparative analysis of the possible further measure, in terms of cost/efficacy, has been carried out sector by sector. In particular, studies showed that a significant margin of reduction can be obtained in the following sectors:

- *industrial sector*. Adoption of best available technologies in the old large plants according to authorisation (environmental permit), as required in 2007 by Annex I of Directive 96/61/EC concerning integrated pollution prevention and control (IPCC).
- *transport sector*. Implementation of technical measures according to Directives on additional measures on light and heavy duty vehicles and non-technical measures, such as rationalization and promotion of public transport for the general population and goods.
- *tertiary industrial and residential sector*. Measures to increase the usage of more efficient household heating plants and appliances and the energy efficiency of buildings; adoption of innovative technologies, usage of low pollution fuels and energy renewable sources.

Results of the studies are reported on the website of the Ministry for the Environment, Land and Sea http://www2.minambiente.it/sito/settori_azione/iar/iam/emissioni/.

References

Amman M., Cofala J., Heyes C., Klimont Z., Shopp W., 1999. The RAINS model: a tool for assessing regional emission control strategies in Europe, Pollution Atmospherique, December 1999.

APAT, 2005. Methodologies used in Italy for the estimation of air emission in the agriculture sector. Technical report 64/2005. Rome - Italy.

URL: http://www.apat.gov.it/site/contentfiles/00140800/140835_R64_2005.pdf.

APAT, 2006. Quality Assurance/Quality Control plan for the Italian Emission Inventory. Procedures Manual. June 2006.

URL: http://www.apat.gov.it/site/it-IT/APAT/Pubblicazioni/Altre_Pubblicazioni.html.

Cóndor R.D., De Lauretis R., 2007. Agriculture emission inventory in Italy: synergies among conventions and directives. In: Ammonia emissions in Agriculture. Ed. G.J. Monteny, E. Hartung. Wageningen Academic Publishers. 404 p. The Netherlands.

Cóndor R.D., Di Cristofaro E., De Lauretis R., 2008. Agricoltura: inventario nazionale delle emissioni e disaggregazione provinciale. Istituto superiore per la protezione e la ricerca ambientale, ISPRA Rapporto tecnico 85/2008. Roma, Italia. URL:

http://www.apat.gov.it/site/it-IT/APAT/Pubblicazioni/Rapporti/Documento/rapporto_85_2008.html

CRPA, 2006[a]. Progetto MeditAIRaneo: settore Agricoltura. Relazione finale. Technical report on the framework of the MeditAIRaneo project for the Agriculture sector, Reggio Emilia - Italy.

CRPA, 2006[b]. Predisposizione di scenari di emissione finalizzati alla progettazione di interventi per la riduzione delle emissioni nazionali di ammoniaca ed alla valutazione di misure e di progetti per la tutela della qualità dell'aria a livello regionale. Final report. Reggio Emilia - Italy.

EEA, 2000. COPERT III, Computer Programme to Calculate Emissions from Road Transport - Methodology and Emission Factors, European Environment Agency, Technical report No 49, November 2000.

EEA, 2007. COPERT 4, Computer programme to calculate emissions from road transport. European Environment Agency, December 2007. URL: <http://lat.eng.auth.gr/copert/docs.htm> .

EMEP/CORINAIR, 2007. Atmospheric Emission Inventory Guidebook. Technical report No 16/2007.

ENEA, 2006. Valutazione della possibilità di sostituzione dell'urea con altri fertilizzanti azotati. Final report. Rome, Italy.

Gracceva F., Contaldi M., 2004. Scenari energetici italiani. Valutazione di misure di politica energetica. ENEA, 2004.

Goldstein, G.A., L.A. Greening, and the Partners in IEA ESAP, 1999. *Energy Planning and the Development of Carbon Mitigation Strategies: Using the MARKAL Family of Models*. White Paper

available from the International Resources Group (IRG). URL: http://www.ecn.nl/unit_bs/etsap .

IIASA, 2008. Technical Documentation on RAINS Europe.
URL: <http://www.iiasa.ac.at/~rains/index.html>.

IPCC, 1997. *Revised 1996 IPCC Guidelines for National Greenhouse Gas Emission Inventories*. Three volumes: Reference Manual, Reporting Manual, Reporting Guidelines and Workbook. IPCC/OECD/IEA. IPCC WG1 Technical Support Unit, Hadley Centre, Meteorological Centre, Meteorological Office, Bracknell, UK.

IPCC, 2000. Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Kanagawa, Japan.

IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan.

ISPRA. 2008[a]. Serie storiche dal 1980 al 2006 delle emissioni nazionali di inquinanti atmosferici, Rete del Sistema Informativo Nazionale Ambientale - SINANET. URL: http://www.sinanet.apat.it/it/sinanet/serie_storiche_emissioni/NFR%201980-2006/view

ISPRA. 2008[b]. Database della disaggregazione a livello provinciale dell'Inventario nazionale delle emissioni:1990-1995-2000-2005. Istituto Superiore per la Protezione e la Ricerca Ambientale, ISPRA. URL: http://www.sinanet.apat.it/it/inventaria/disaggregazione_prov2005/

ISPRA, 2009 [a]. National Greenhouse Gas Inventory System in Italy. April 2009.
URL: http://www.apat.gov.it/site/it-IT/APAT/Pubblicazioni/Altre_Pubblicazioni.html

ISPRA, 2009 [b]. Italian Greenhouse Gas Inventory 1990-2007. National Inventory Report 2009. April 2009.
URL:http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/4771.php.

ISPRA, 2009 [c]. Quality Assurance/Quality Control plan for the Italian Emission Inventory. Year 2009. April 2009.
URL: http://www.apat.gov.it/site/it-IT/APAT/Pubblicazioni/Altre_Pubblicazioni.html.

ISPRA, 2009 [d]. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009.

ISPRA, 2009[e]. Annuario dei dati ambientali 2008. Capitolo ATMOSFERA. Roma, Italia.
URL: http://annuario.apat.it/capitoli/Ver_6/versione_integrale/06%20Atmosfera.pdf .

MATTM, 2007. Fourth National Communication under the UN Framework Convention on Climate Change. Italy. Ministry for the Environment, Land and Sea. November 2007.

Pignatelli T., De Lauretis R., Contaldi M., D'Elia I., Romano D., Vialetto G., 2007. Harmonization of National inventory and projections of multi-pollutant emission scenarios. The Italian experience within the European context and the UN-ECE Convention on Long Range Transboundary Air Pollution. 16th Annual International Emission Inventory Conference USEPA .

Romano D., Bernetti A., De Lauretis R., 2004. Different methodologies to quantify uncertainties of air emissions. Environment International vol 30 pp 1099-1107.

Techne, 2009. Stima delle emissioni in atmosfera nel settore del trasporto aereo e marittimo. Final report. TECHNE Consulting, March 2009.

UNECE, 2003. Guidelines for Estimating and Reporting Emission Data under the Convention on Long-range Transboundary Air Pollution, Emission Reporting Guidelines. Air Pollution Studies no. 15, 2003. URL: <http://www.unece.org/env/documents/2003/eb/air/ece.eb.air.80.E.pdf>.

Vialetto G., Contaldi M., De Lauretis R., Lelli M., Mazzotta V., Pignatelli T., 2005. Emission Scenarios of Air Pollutants in Italy using Integrated Assessment Models. Pollution Atmosphérique, vol. 185 pp.71-78.

Zanini G., Pignatelli T., Monteforti F., Vialetto G., Vitali L., Brusasca G., Calori G., Finardi S., Radice P., Silibello C., 2005. The MINNI Project: An Integrated Assessment Modelling System for Policy making. In Zenger A. and Argent R.M. (eds) MODSIM 2005 International Congress on Modelling and Simulation. Modelling and Simulation Society of Australia and New Zealand, December 2005, pp. 2005-2011. ISBN: 0-9758400-2-9.
URL: <http://www.mssanz.org.au/modsim05/papers/zanini.pdf>.

Annex: NFR codes

1 A 1	Energy industries
1 A 1 a	Public Electricity and Heat Production
1 A 1 b	Petroleum refining
1 A 1 c	Manufacture of Solid Fuels and Other Energy Industries
1 A 2	Manufacturing Industries and Construction
1 A 2 a	Iron and Steel
1 A 2 b	Non-ferrous Metals
1 A 2 c	Chemicals
1 A 2 d	Pulp, Paper and Print
1 A 2 e	Food Processing, Beverages and Tobacco
1 A 2 f	Other
1 A 3	Transport
1 A 3 a ii (i)	Civil Aviation (Domestic, LTO)
1 A 3 a i (i)	Civil Aviation (International, LTO)
1 A 3 b	Road Transportation
1 A 3 b i	R.T., Passenger cars
1 A 3 b ii	R.T., Light duty vehicles
1 A 3 b iii	R.T., Heavy duty vehicles
1 A 3 b iv	R.T., Mopeds & Motorcycles
1 A 3 b v	R.T., Gasoline evaporation
1 A 3 b vi	R.T., Automobile tyre and brake wear
1 A 3 b vii	R.T., Automobile road abrasion
1 A 3 c	Railways
1 A 3 d ii	National Navigation
1 A 3 e	Other
1 A 3 e i	Pipeline compressors
1 A 3 e ii	Other mobile sources and machinery
1 A 4	Other Sectors
1 A 4 a	Commercial / Institutional
1 A 4 b	Residential
1 A 4 b i	Residential plants
1 A 4 b ii	Household and gardening (mobile)
1 A 4 c	Agriculture / Forestry / Fishing
1 A 4 c i	Stationary
1 A 4 c ii	Off-road Vehicles and Other Machinery

1A 4 c iii	National Fishing
1 A 5 a	Other, Stationary (including Military)
1 A 5 b	Other, Mobile (Including military)
1B 1	Fugitive Emissions from Solid Fuels
1 B 1 a	Coal Mining and Handling
1 B 1 b	Solid fuel transformation
1 B 1 c	Other
1 B 2	Oil and natural gas
1 B 2 a	Oil
1 B 2 a i	Exploration Production, Transport
1 B 2 a iv	Refining / Storage
1 B 2 a v	Distribution of oil products
1 B 2 a vi	Other
1 B 2 b	Natural gas
1 B 2 c	Venting and flaring
2 A	Mineral Products
2 A 1	<i>Cement Production</i>
2 A 2	<i>Lime Production</i>
2 A 3	<i>Limestone and Dolomite Use</i>
2 A 4	<i>Soda Ash Production and use</i>
2 A 5	<i>Asphalt Roofing</i>
2 A 6	<i>Road Paving with Asphalt</i>
2 A 7	<i>Other including Non Fuel Mining & Construction</i>
2 B	Chemical industry
2 B 1	<i>Ammonia Production</i>
2 B 2	<i>Nitric Acid Production</i>
2 B 3	<i>Adipic Acid Production</i>
2 B 4	<i>Carbide Production</i>
2 B 5	<i>Other</i>
2 C	Metal Production
2 D	Other Production
2 D 1	<i>Pulp and Paper</i>
2 D 2	<i>Food and Drink</i>
2 G	Other
3 A	Paint Application
3 B	Degreasing and dry cleaning
3 C	Chemical products, manufacture and processing

3 D	Other including products containing HMs and POPs
4 B	Manure management
4 B 1	<i>Cattle</i>
4 B 1 a	<i>Dairy</i>
4 B 1 b	<i>Non-Dairy</i>
4 B 2	<i>Buffalo</i>
4 B 3	<i>Sheep</i>
4 B 4	<i>Goats</i>
4 B 5	<i>Camels and Llamas</i>
4 B 6	<i>Horses</i>
4 B 7	<i>Mules and Asses</i>
4 B 8	<i>Swine</i>
4 B 9	<i>Poultry</i>
4 B 13	<i>Other</i>
4 C	Rice cultivation
4 D 1	<i>Direct Soil Emission</i>
4 F	Field burning of agricultural wastes
4 G	Other
5 B	Forest and grassland conversion
6 A	Solid waste disposal on land
6 B	Waste-water handling
6 C	Waste incinerator
6 D	Other waste